

CONTRIBUTIONS
FROM THE
CUSHMAN LABORATORY
FOR
FORAMINIFERAL RESEARCH

INDEX TO VOLUMES 21 — 25

PRICE \$1.00 POSTPAID

SHARON, MASSACHUSETTS, U. S. A.

CONTRIBUTIONS FROM THE CUSHMAN LABORATORY FOR FORAMINIFERAL RESEARCH

INDEX TO VOLUMES 21-25 INCLUSIVE

Owing to the demand of those who wish to bind these Contributions, the following Index has been prepared.

All new genera and species appear in **bold-faced type**, others in *italic type*. An abbreviated form gives the geologic formation and general locality with each species.

Papers are grouped both by geographic and geologic headings as well as by subjects.

In order to conserve space, the following abbreviated form has been used: Volume number with the part number following in parentheses followed by a colon. After the colon comes the page number, then the plate number in **bold-faced type**, and the figure number in *italics*. It is hoped that this will give ready reference to the subject matter of these volumes.

Additional Homonyms in Foraminifera Erected Since 1940, 22(4) : 123.

Additional New Foraminifera from the American Paleocene, 24(2) : 43.

Additional New Species of Arenaceous Foraminifera from Shallow Waters of Trinidad, 24(2) : 37.

Additional Species of Paleocene Foraminifera from the Madruga Formation of Cuba, 24(4) : 85.

[Alabama]. A Foraminiferal Fauna from the Lisbon Formation of Alabama, 21(1) : 11. A New *Pseudoclavulina* from the Upper Cretaceous of Alabama, 21(3) : 74. Some Foraminifera of Woodbine Age from Texas, Mississippi, Alabama, and Georgia, 22(3) : 71.

Alabamina, Some Notes on the Genera *Pulvinulinella*, *Parrella*, and, 24(1) : 13.

Alabamina, 24(1) : 14; *wilcoxensis* (Olig.-Miss.), 24(1) : 10; (Eoc.-Ala.), 14, 2, 18-20; (Eoc.-Miss.), 24(2) : 35, 6, 28, 29.

[Alaska]. A Foraminiferal Fauna from Amchitka Island, Alaska, 23(3) : 60.

[Algeria]. *Globigerinelloides*, A New Genus of the Globigerinidae, 24(2) : 42.

Allomorphina and *Quadrिमorphina*, Species of the Genera, 25(3) : 59.

Allomorphina, 25(3) : 59; *abbreviata* (Eoc.-France), 25(3) : 66, 11, 17; *conica* (Cret.-Amer.), 25(3) : 62, 11, 8; *cretacea* (Cret.-Europe, Amer.), 25(3) : 61, 11, 3, 4; *cubensis* (Olig.-Cuba), 25(3) : 67, 11, 19, 20; *globulosa* (Paleoc.-Tex.), 25(3) : 64, 11, 14; *halli* (Eoc.-N. J.), 25(3) : 66, 11, 18; *macrostoma* (Mioc.-Rec.), 25(3) : 68, 12, 4, 5; *minuta* (Cret.-Tex.), 25(3) : 62, 11, 7; *navarroana* (Cret.-Tex.), 25(3) : 61, 11, 6; *obliqua* (Cret.-Europe), 25(3) : 60, 11, 2; *pacifica* (Mioc. to Rec.-Indo-Pac.), 25(3) : 68, 12, 6-9; *paleocenica* (Paleoc.-Tex.), 24(2) : 45, 8, 10; (Paleoc.-Tex., Ala., Miss., Ark.), 25(3) : 65, 11, 16; *prima* (Jur.-France), 25(3) : 60, 11, 1; *subtriangularis* (Paleoc.-Miss.), 25(3) : 64, 11, 15; *trigona* (Paleoc.-Ark.), 22(2) : 63, 11, 11, 15; Mioc., Olig.-Europe), 25(3) : 67, 12, 1, 2; var. *obtusa* (Olig.-France), 25(3) : 67, 12, 3; *trochoides* (Cret.-Trinidad), 23(2) : 49; *velascoensis* (Cret.-Trinidad), 23(2) : 49; (Cret.-Mex.), 25(3) : 61, 11, 5; *whangaia* (Cret.-N. Zealand), 25(3) : 64, 11, 9-13.

Allomorphinella, 25(4) : 98; *contraria* (Cret.-Europe), 25(4) : 98, 16, 28.

Amendment to the Diagnosis of *Lockhartia huntii* Ovey, 24(4) : 99.

[America]. Some New Foraminifera from the American Cretaceous, 23(3) : 53. Some New Foraminifera from the Paleocene of the Southern United States, 23(4) : 81. Additional New Foraminifera from the American Paleocene, 24(2) : 43.

Ammoastuta, 24(1) : 17; *salsa* (Rec.-Trinidad), 24(1) : 17, 3, 14-16.

Ammobaculites paleocenicus Cushman, A New Name, 23(4) : 77.

Ammobaculites advenus (Cret.-Tex.), 23(3) : 53, 13, 1; *agrestis* (Cret.-Tex., Fla., Ga., Ala.), 23(3) : 53, 13, 2, 3; *alexanderi* (Cret.-Trinidad), 23(2) : 37, 11, 6; *bergquisti* (Cret.-Tex., Ala., Ga.), 22(3) : 73, 13, 4, 5; *braunsteini* (Cret.-Miss., Tex., Ala.), 22(3) : 74, 13, 7; *comprimatus* (Cret.-Tex.), 22(3) : 73, 13, 3; *coprolithiformis* (Cret.-Trinidad), 23(2) : 37; *dilatatus* (Rec.-Trinidad), 24(2) : 39, 7, 10, 11; *directus* (Rec.-Trinidad), 24(2) :

38, 7, 3, 4; *diversus* (Rec.-Trinidad), 24(2) : 38, 7, 5, 6; *exiguus* (Rec.-Trinidad), 24(2) : 38, 7, 7, 8; *exilis* (Rec.-Trinidad), 24(2) : 39, 7, 9; *foliaceus* (Eoc.-Peru), 25(4) : 76, 13, 14; *fragmentarius* (Cret.-Tex.), 22(3) : 74; *gratus* (Cret.-Tex., Fla., Ga., Ala.), 23(3) : 54, 13, 4; *jarvisi* (Cret.-Venezuela), 23(1) : 4, 1, 10-12; (Cret.-Trinidad), 23(2) : 37; *junceus* (Cret.-Tex.), 22(3) : 72, 13, 2; *midwayensis* (Paleoc.-Ala.), 23(4) : 77; *paleocenicus* (Paleoc.-Ala.), 23(4) : 77; *pseudocassis* (Rec.-Trinidad), 24(2) : 39, 7, 12; *salsus* (Rec.-Trinidad), 24(1) : 16, 3, 7-9; var. *distinctus* (Rec.-Trinidad), 24(2) : 40, 7, 13, 14; *stephensoni* (Cret.-Tex., Ala., Ga.), 22(3) : 72, 13, 1; sp. (Cret.-Venezuela), 23(1) : 4, 1, 9.

Ammobaculoides plummerae (Cret.-Tex., Ga.), 22(3) : 74, 13, 6.

Ammodiscus asper (Jur.-Poland), 25(1) : 3, 1, 1; *cretaceus*, var. *rugosa*, 23(4) : 92; *glabratus* (Cret.-Venezuela), 23(1) : 3, 1, 3; (Cret.-Trinidad), 23(2) : 35; (Eoc.-Peru), 25(4) : 75, 13, 9; *gullmarensis* (Rec.-Atl.), 24(2) : 45; *incertus* (Eoc.-Peru), 25(4) : 75, 13, 8; *pennyi* (Cret.-Venezuela), 23(1) : 3, 1, 4; (Cret.-Trinidad), 23(2) : 35; *planus* (Rec.-Atl.), 24(2) : 45; *restinensis* (Eoc.-Peru), 25(3) : 50, 9, 3.

Ammolagena clavata (Cret.-Trinidad), 23(2) : 36.

Ammospirata mexicana, 21(2) : 49.

Amphimorphina fragilis (Eoc.-Miss.), 24(1) : 10, 2, 1, 2; 24(2) : 46; *gracilis* (Eoc.-Miss.), 24(1) : 3, 2, 1, 2; 24(2) : 46.

Amphistegina from the Eocene of Ecuador, A New Species of, 22(4) : 117.

Amphistegina elliotti (Eoc.-Ecuador), 22(4) : 118, 20, 1-6.

Angulogerina byramensis (Olig.-Miss.), 22(3) : 95; 24(1) : 9; *charapotoensis* (Mioc.-Ecuador), 24(3) : 63, 10, 8; *cooperensis* (Eoc.-Ga.), 21(3) : 66, 10, 28; (Eoc.-Miss.), 21(4) : 99, 15, 24; *fluens* (Plioc.-Alaska), 23(3) : 67, 16, 6, 7; *mauricensis* (Eoc.-Miss.), 24(2) : 33, 6, 11; *ocalana* (Eoc.-Ga.), 21(1) : 8, 2, 9; (Eoc.-Miss.), 21(4) : 99, 15, 23; *rugoplicata* (Olig.-Miss.), 22(3) : 95; *vicksburgensis* (Eoc.-Ga.), 21(3) : 67, 10, 27; (Olig.-Miss.), 22(3) : 95; 24(1) : 9; *wilcoxensis* (Paleoc.-Cuba), 24(4) : 89.

Anomalina tenuissima (Reuss), var. *evoluta* van Bellen, a Homonym, 24(3) : 76.

Anomalina acuta (Paleoc.-Ark.), 22(2) : 64, 11, 13, 14; (Paleoc.-Cuba), 24(4) : 89; *beccariiformis* (Cret.-Trinidad), 23(2) : 50, 12, 14; *bengalensis* (Plioc.-Kar Nicobar), 24(1) : 13, 2, 17; *bi-*

- lateralis* (Eoc.-Miss.), 21(4): 103, 16, 14, 15; (Olig.-Miss.), 24(1): 10; *clementiana* (Paleoc.-Ark.), 22(2): 64, 11, 16, 17; var. *assimilis* (Paleoc.-Cuba), 24(3): 74, 12, 17; *costiana* (Eoc.-Ga.), 21(3): 72; *cubana* (Paleoc.-Cuba), 24(4): 86, 15, 7-9; *henbesti* (Cret.-Venezuela), 23(1): 18, 4, 32; *madrugensis* (Paleoc.-Cuba), 24(4): 86, 15, 4-6; *martinezensis* (Paleoc.-Cuba), 24(3): 74, 12, 14-16; *midwayensis* (Paleoc.-Ark.), 22(2): 64, 11, 18, 19; *obesa* (Cret.-Miss.), 23(3): 55, 13, 7; *polyrraphes* (Cret.-Trinidad), 23(2): 51; *praespissiformis* (Paleoc.-Cuba), 24(4): 86, 15, 1-3; *reinholdi* (Plioc.-Holland), 23(3): 59, 14, 7; *restinensis* (Eoc.-Peru), 25(3): 57, 10, 17; *rubiginosa* (Cret.-Venezuela), 23(1): 18, 4, 33; (Cret.-Trinidad), 23(2): 50; *tenuissima*, var. *evoluta* (Eoc.-Dalmatia), 24(3): 76; var. *jongmansi* (Eoc.-Dalmatia), 24(3): 76; *variolata*, var. *evoluta* (Rec.-Italy), 24(3): 76; sp. (Cret.-Venezuela), 23(1): 18, 4, 30.
- Arenobulimina americana* (Cret.-Venezuela), 23(1): 7, 2, 9; (Cret.-Trinidad), 23(2): 38, 11, 7; *conica*, 22(4): 131; *conoidea*, 22(4): 131; *jurassica* (Jur.-Poland), 25(1): 8, 2, 17, 18; sp. (Cret.-Calif.), 24(4): 93, 16, 10.
- [Arkansas]. A Foraminiferal Fauna from the Paleocene of Arkansas, 22(2): 45. Two New Species of Foraminifera from the Arkadelphia Marl of Arkansas, 23(3): 56.
- Articulina advena* (Olig.-Miss.), 22(3): 81, 14, 8, 9; 24(1): 8; *atlantica* (Rec.-Atl.), 23(4): 89, 19, 6; *byramensis* (Olig.-Miss.), 22(3): 82, 14, 10, 11; 24(1): 8; *jacksonensis* (Olig.-Miss.), 24(1): 8, 1, 4; *terquemi* (Eoc.-Miss.), 21(4): 83, 13, 20; sp. (Olig.-Miss.), 22(3): 82, 14, 13.
- Assilina sublacunata*, 23(4): 92; *subumbilicata*, 23(4): 93.
- Asterigerina byramensis* (Olig.-Miss.), 22(3): 101, 16, 28, 29; *crassaformis* (Eoc.-Peru), 25(4): 82, 14, 16; *lisbonensis* (Eoc.-Ala.), 21(1): 19, 4, 15-18; *subacuta* (Olig.-Miss.), 22(3): 101, 16, 27.
- Astronion viragoense* (Plioc.-Alaska), 23(3): 64.
- Astrorhiza vermiformis* (Eoc.-Peru), 25(4): 74, 13, 1.
- [Atlantic]. A New Species of *Globobulimina* from the Western Atlantic, 21(3): 73. New Species and Varieties of Foraminifera from off the Southeastern Coast of the United States, 23(4): 86. New Names for Four Homonym Species Described in "Foraminifera in the Gullmar Fjord and the Skagerak," 24(2): 45.

Bathysiphon alexanderi (Cret.-Trinidad), 23(2) : 34, 11, 1; *dubia* (Cret.-Trinidad), 23(2) : 33; *eocenica* (Eoc.-Peru), 25(4) : 75, 13, 3.

Bifarina vicksburgensis (Olig.-Miss.), 22(3) : 94, 16, 1; 24(1) : 9.
Bitubulogenerina aperta (Olig.-Miss.), 22(3) : 94, 16, 2; *ellisi* (Eoc.-Ala.), 21(1) : 18, 4, 10; *howei* (Olig.-Miss.), 24(1) : 9; *moodysensis* (Eoc.-Miss.), 21(4) : 98, 15, 25, 26; *vicksburgensis* (Olig.-Miss.), 22(3) : 94, 16, 3; 24(1) : 9.

Boldia, 22(4) : 122; *carinata* (Paleoc.-Cuba), 24(3) : 75, 11, 11-13; *cubensis* (Paleoc.-Cuba), 24(3) : 74, 11, 15, 16; *lobata* (Eoc.-France), 22(4) : 122; *madruagaensis* (Paleoc.-Cuba), 24(3) : 75, 12, 4-6; *vandersluisi* (Eoc.-Guatemala & Brit. Honduras), 22(4) : 122, text fig. 1.

Bolivina acerosa, var. *pacifica*, 25(4) : 101; *alata* (Mioc.-Ecuador), 24(3) : 60, 9, 32; *alazanensis* (Eoc.-Ga.), 21(3) : 66; (Eoc.-Miss.), 21(4) : 97, 15, 22; *astoriensis*, 25(4) : 100; *broussardi* (Eoc.-Ga.), 21(3) : 66, 10, 24; *byramensis* (Olig.-Miss.), 22(3) : 92, 15, 29; *charapotoensis* (Mioc.-Ecuador), 24(3) : 61, 10, 1, 2; *chehalisensis*, 25(4) : 100; *choctawensis* (Olig.-Miss.), 22(3) : 93, 15, 32; *cuneiformis* (Mioc.-Ecuador), 24(3) : 60, 9, 31; *dispar* (Mioc.-Ecuador), 24(3) : 60, 9, 35; *dissentiata* (Mioc.-Jamaica), 22(4) : 120; *ecuadorana* (Mioc.-Ecuador), 24(3) : 60, 9, 33, 34; *gracilis* (Eoc.-Ga.), 21(3) : 66, 10, 23; (Eoc.-Miss.), 21(4) : 96, 15, 17; 24(1) : 10; var. *danvillensis* (Eoc.-Miss.), 21(4) : 96, 15, 18; *granti* (Mioc.-Ecuador), 24(3) : 59, 9, 36; var. *denticulata* (Mioc.-Ecuador), 24(3) : 59, 9, 37; *ignara* (Eoc.-Peru), 25(3) : 54, 10, 3, 4; *imporcata* (Mioc.-Jamaica), 22(4) : 120; *incrassata* (Cret.-Venezuela), 23(1) : 15, 4, 20; *insueta* (Eoc.-Peru), 25(3) : 53, 10, 1, 2; *interjuncta*, var. *bicostata* (Mioc.-Ecuador), 24(3) : 59, 9, 29, 30; *irregularis*, 22(4) : 130; *jacksonensis* (Eoc.-Ga.), 21(1) : 8; 21(3) : 66; (Eoc.-Miss.), 21(4) : 95, 15, 14; (Olig.-Miss.), 22(3) : 93; var. *striatella* (Eoc.-Ga.), 21(1) : 8; (Eoc.-Miss.), 21(4) : 96, 15, 15; *louisiana* (Eoc.-Miss.), 21(4) : 96, 15, 16; *mauricensis* (Eoc.-Miss.), 24(2) : 33, 6, 9; *mississippiensis* (Olig.-Miss.), 22(3) : 92, 15, 30; var. *costifera* (Olig.-Miss.), 22(3) : 93, 15, 31; 24(1) : 9; *moodysensis* (Eoc.-Miss.), 21(4) : 97, 15, 19, 20; *mornhinvegi* (Olig.-Miss.), 22(3) : 93, 15, 33; *pseudoplicata* (Plioc.-Alaska), 23(3) : 66, 16, 2, 3; *robusta*, var. *pacifica*, 25(4) : 101; *scintillo* (Eoc.-Miss.), 24(1) : 4, 2, 5; 10, 2, 5; *subangularis*, var. *irregularis*, 22(4) : 130; 23(4) : 93; *taylori* (Eoc.-Ala.), 21(1) : 17, 4,

- 9; (Eoc.-Miss.), 21(4) : 97, 15, 21; 24(2) : 33, 6, 10; sp. (Cret.-Calif.), 24(4) : 95, 16, 14.
- Bolivinella*, 22(4) : 131; *rugosa* (Olig.-Miss.), 24(1) : 8; *subpectinata* (Olig.-Miss.), 22(3) : 91; var. *interrupta* (Olig.-Miss.), 22(3) : 91; *vicksburgensis* (Olig.-Miss.), 24(1) : 8.
- Bolivinella eleyi* (Cret.-Trinidad), 23(2) : 44, 11, 12.
- Bolivinella*, 22(4) : 131.
- Bolivinoides trinitatensis* (Cret.-Trinidad), 23(2) : 44.
- Bolivinopsis pulchella* (Eoc.-Ecuador), 23(4) : 78, 17, 4, 5.
- Bulimina* from the Eocene, Chira Shale, of Peru, A New Species of, 23(3) : 57.
- Bulimina acutangularis* (Eoc.-Ecuador), 23(4) : 80, 17, 13; *arkadelphiana*, var. *midwayensis* (Paleoc.-Cuba), 24(4) : 89; *aspera* (Cret.-Venezuela), 23(1) : 14, 4, 17; *byramensis* (Olig.-Miss.), 22(3) : 91, 15, 25, 26; *cacumenata* (Paleoc.-Ark.), 22(2) : 59, 10, 20; *cooperensis* (Eoc.-Ga.), 21(1) : 8; *decurtata* (Eoc.-Ecuador), 23(4) : 80, 17, 11; *ecuadorana* (Mioc.-Ecuador), 24(3) : 58, 9, 25, 26; *elongata*, var. *tenera* (Eoc.-Ga.), 21(3) : 64, 10, 16; *jacksonensis* (Eoc.-Miss.), 24(1) : 10; var. *cuneata* (Eoc.-Miss.), 24(1) : 10; *jugosa* (Mioc.-Ecuador), 24(3) : 57, 9, 22; *kickapoensis*, var. *pingua* (Cret.-Trinidad), 23(2) : 45, 12, 3; *limbata* (Cret.-Trinidad), 23(2) : 45, 12, 4; *lineata* (Eoc.-Ecuador), 23(4) : 79, 17, 12; *pagoda*, var. *hebespinata* (Mioc.-Ecuador), 24(3) : 58, 9, 23; *petroleana* (Cret.-Venezuela), 23(1) : 14, 4, 18, 19; (Cret.-Trinidad), 23(2) : 45; var. *spinea* (Cret.-Trinidad), 23(2) : 45; *punctato-costata* (Eoc.-Peru), 25(3) : 52, 9, 12, 13; *pupoides* (Eoc.-Peru), 25(4) : 79, 14, 6, 7; *secaensis* (Eoc.-Ecuador), 23(4) : 79, 17, 9, 10; *simplex* (Eoc.-Ala.), 21(1) : 17, 4, 5; (Eoc.-Miss.), 24(2) : 32, 6, 7; *trigona* (Eoc.-Ala.), 21(1) : 17, 4, 6; 22(4) : 131; *trigonalis*, 22(4) : 131; *trigonula*, 22(4) : 131; *uvigeriniformis*, var. *charapotoensis* (Mioc.-Ecuador), 24(3) : 58, 9, 24; sp. (Cret.-Calif.), 24(4) : 94, 16, 13.
- Bulimina (Desinobulimina) auriculata* (Olig.-Miss.), 22(3) : 92; *diversa* (Eoc.-Peru), 23(3) : 57, 14, 1-3; *expansa* (Eoc.-Peru), 25(3) : 52, 9, 14; *quadrata* (Paleoc.-Ark.), 22(2) : 60, 10, 19; (Paleoc.-Cuba), 24(4) : 89; *suteri* (Cret.-Venezuela), 23(1) : 14, 4, 16.
- Buliminella carseyae*, var. *plana* (Cret.-Trinidad), 23(2) : 45, 12, 1; *colonensis* (Cret.-Trinidad), 23(2) : 45, 12, 2; *curta* (Mioc.-Ecuador), 24(3) : 56, 9, 21; *ecuadorana* (Mioc.-Ecuador),

24(3): 57, 9, 19, 20; *elegantissima* (Eoc.-Ga.), 21(1): 7, 2, 6; (Plioc.-Alaska), 23(3): 65, 15, 21; *madagascariensis*, var. *spicata* (Olig.-Miss.), 22(3): 91, 15, 23, 24; 24(1): 9; *obtusata* (Olig.-Miss.), 22(3): 91, 15, 21, 22; *robertsi* (Eoc.-Ga.), 21(3): 64, 10, 15; (Eoc.-Miss.), 21(4): 94, 15, 12.

Buliminellita, 23(4): 78; *mirifica* (Eoc.-Ecuador), 23(4): 78, 17, 6-8.

Buliminidae, The Species of the Subfamily Reussellinae of the Foraminiferal Family, 21(2): 23.

Bullopore chapmani (Paleoc.-Ark.), 22(2): 58, 10, 13, 14; (Paleoc.-Cuba), 24(3): 71, 11, 8; var. *hispida*, 22(4): 130; *laevis*, var. *hispida*, 22(4): 129.

[California]. Some New Species and Varieties of Foraminifera from the Pliocene of Timms Point, California, 22(2): 65. A Foraminiferal Fauna from the New Almaden District, California, 24(4): 90.

Camerina jacksonensis (Eoc.-Miss.), 21(4): 93, 14, 22, 23.

Cancris claibornensis (Eoc.-Ala.), 21(1): 19; (Eoc.-Miss.), 24(2): 35, 6, 19; *cocoaensis* (Eoc.-Miss.), 24(1): 11; *cubensis* (Eoc.-Ga.), 21(3): 69, 11, 7; *pauciloculatus* (Olig.-Miss.), 22(3): 100.

Candorbulina universa (Mioc.-Ecuador), 24(3): 66, 10, 23.

Cassidulina californica (Plioc.-Alaska), 23(3): 68, 16, 16-18; *crassa* (Olig.-Miss.), 24(1): 10; *delicata* (Mioc.-Ecuador), 24(3): 65, 10, 20; *diversa* (Eoc.-Peru), 25(3): 56, 10, 14; *globosa* (Eoc.-Ga.), 21(3): 70; (Eoc., Olig.-Miss.), 24(1): 11; (Eoc.-Peru), 25(4): 82, 14, 14; *inconspicua*, 25(4): 101; *insueta* (Rec.-Atl.), 23(4): 92, 20, 10; *limbata* (Plioc.-Alaska), 23(3): 68, 16, 22; *lomitensis* (Plioc.-Alaska), 23(3): 69, 16, 19; (Mioc.-Ecuador), 24(3): 65, 10, 19; *moodyensis* (Eoc.-Miss.), 21(4): 102, 16, 9, 10; (Olig.-Miss.) 22(3): 101; *murrhyna*, 25(1): 21; *parva*, 25(4): 101; *quadrata*, var. (Plioc.-Alaska), 23(3): 69, 16, 20, 21; *subglobosa*, var. *parva*, 25(4): 101; *twiggsana* (Eoc.-Ga.), 21(1): 10, 2, 13; *winniana* (Eoc.-Ga.), 21(3): 70, 11, 10.

Cassidulinoides bradyi (Plioc.-Alaska), 23(3): 69, 16, 23.

Ceratobulimina and Its Species, The Genus, 22(4): 107.

Ceratobulimina, 22(4): 107; *alazanensis* (Olig.-Mex.), 22(4): 111, 18, 17, 18; *bundensis* (Eoc.-Holland), 22(4): 109; *carpatica* (Mioc.-Galicia), 22(4): 113, 18, 7; *chapmani* (Rec.-Antarctic),

22(4) : 114, 19, 2-4; *contraria* (Olig.-Mioc.), 22(4) : 110, 17, 10, 11; *cretacea* (Cret.-Tex., Ark., Tenn., Miss.), 22(4) : 108, 17, 1, 2; *dehiscens* (Mioc.-Australia), 22(4) : 112, 18, 8-10; *evoluta* (Olig.-Trinidad), 22(4) : 112, 18, 4; *eximia* (Eoc.-Amer., Europe), 22(4) : 109, 17, 6-8; *inflata* (Eoc.-Holland), 22(4) : 117, 19, 10; *kellumi* (Mioc.-New Zealand), 22(4) : 113, 18, 6; *lenticula* (Cret.-France), 22(4) : 117; *lornensis* (Olig.-N. Zealand), 22(4) : 112, 18, 5; *pacifica* (Mioc. to Rec.-Pac.), 22(4) : 113, 18, 11-16; *perplexa* (Paleoc., Eoc.-Amer. Europe), 22(4) : 108, 17, 3-5; *tenuis* (Rec.-Antarctic), 22(4) : 114, 19, 1; *washburnei* (Olig.-Ore., Wash.), 22(4) : 111, 18, 1-3; *westraliensis* (Eoc.-Australia), 22(4) : 110, 17, 9.

Ceratobulimina (*Ceratocancris*) *australis* (Tert.-Australia), 22(4) : 116, 19, 8, 9; *clifdenensis* (Mioc.-N. Zealand), 22(4) : 115, 19, 7; *hauerii* (Mioc.-Austria), 22(4) : 115, 19, 5, 6.

Ceratocancris, 22(4) : 115.

Cercidina, 22(4) : 131.

Chilostomella and Related Genera, Species of the Genus, 25(4) : 84.

Chilostomella, 25(4) : 85; *chilostomelloides* (Eoc.-Moravia), 25(4) : 87, 15, 7; *cushmani* (Rec.-Pac.), 25(4) : 93, 16, 4, 5, 11, 12; *cylindroides* (Tert. & Rec.), 25(4) : 87, 15, 8-10; var. *tenuis* (Olig., Eoc.-Europe), 25(4) : 88, 15, 11; *czizeki* (Olig.-Rec.), 25(4) : 90, 15, 20-22; *globata* (Tert.-Porto Rico), 25(4) : 89, 15, 15; *grandis* (Rec.-Pac.), 25(4) : 92, 16, 1, 2, 6; *hadleyi* (Tert.-Cuba), 25(4) : 88, 15, 12-14; *mediterraneensis* (Rec.-Medit.), 25(4) : 92, 15, 25, 26; *mexicana* (Eoc.-Mex.), 25(4) : 86, 15, 3, 4; var. *chirana* (Eoc.-Peru), 25(4) : 86, 15, 6; var. *serrata* (Eoc.-Peru), 25(4) : 86, 15, 5; *milletti* (Rec.-Pac.), 25(4) : 93, 16, 3; *oolina* (Mioc.-Rec.), 25(4) : 91, 15, 23, 24; (Tert.-Borneo), 25(4) : 96; *ovoidea* (Cret.-Venezuela), 23(1) : 16, 4, 28; (Mioc.-Austria; Rec.-Pac.), 25(4) : 89, 15, 17-19; *primitiva* (Cret.-Venezuela), 25(4) : 85, 15, 1; *trinitatensis* (Cret.-Trinidad), 25(4) : 85, 15, 2; *urceolus* (Tert.-Porto Rico), 25(4) : 89, 15, 16.

Chilostomellina, 25(4) : 99, *fimbriata* (Rec.-Pac.), 25(4) : 99, 16, 29.

Chilostomelloides, 25(4) : 93; *cyclostoma* (Tert.-Europe), 25(4) : 96, 16, 17; *eocenica* (Eoc.-Amer.), 25(4) : 94, 16, 10; *ovicula* (Eoc.-Peru), 25(4) : 82, 14, 20; (Tert.-Trinidad), 25(4) : 95, 16, 15, 16; *oviformis* (Eoc., Olig.-Europe, Amer.), 25(4) : 94, 16,

7-9; n. sp. (Eoc.-Dalmatia), 25(4) : 95, 16, 13, 14; sp. (Cret.-Peru), 25(4) : 95.

Chrysalidinella, 21(2) : 49; *costata* (Mioc.-Australia), 21(2) : 52, 8, 19, 20; *cubana* (Eoc.-Cuba), 21(2) : 50, 8, 13-15; *cubensis* (Olig.-Cuba), 21(2) : 50, 8, 16; *dimorpha* (Rec.-Pac.), 21(2) : 52, 8, 21, 22; *earlandi* (Rec.-Pac.), 21(2) : 53, 8, 23; *fijiensis* (Rec.-Fiji), 21(2) : 53, 8, 24; *miocenica* (Mioc.-Fla.), 21(2) : 51, 8, 18; *pulchella* (Mioc.-Panama), 21(2) : 51, 8, 17.

Chrysalogonium arkansasanum (Paleoc.-Ark.), 22(2) : 52, 9, 1, 2; *atlanticum* (Rec.-Atl.), 23(4) : 90, 19, 10; *cretaceum* (Cret.-Venezuela), 23(1) : 12, 3, 27; *eocenicum* (Paleoc.-Ark.), 22(2) : 53, 9, 3-5; *velascoense* (Cret.-Trinidad), 23(2) : 42; sp. (Olig.-Miss.), 22(3) : 86, 15, 1, 2.

Cibicides alaskensis (Plioc.-Alaska), 23(3) : 72, 16, 31, 32; *alleni* (Paleoc.-Cuba), 24(4) : 89; *americanus* (Olig.-Miss.), 22(3) : 102, var. *antiquus* (Eoc.-Miss.), 21(4) : 104, 16, 21, 22; *beaumontianus* (Cret.-Venezuela), 23(1) : 18, 4, 35; *blanpiedi* (Paleoc.-Cuba), 24(4) : 89; *choctawensis*, var. *dorso-involuta* (Eoc.-Holland), 22(4) : 122; var. *ornata* (Eoc.-Holland), 22(4) : 121; *constrictus* (Cret.-Trinidad), 23(2) : 51; *cooperensis* (Eoc.-Miss.), 24(1) : 11, 2, 11; *cushmani* (Eoc.-Peru), 25(4) : 83, 14, 26; *danvillensis* (Eoc.-Ga.), 21(3) : 72, 11, 14; *grimsdalei* (Eoc.-Peru), 25(4) : 83, 14, 24; *harperi* (Cret.-Venezuela), 23(1) : 18, 4, 34; *howei* (Eoc.-Ala), 21(1) : 20, 4, 21; *isidroensis* (Mioc.-Ecuador), 24(3) : 67, 10, 30, 31; *lalickeri* (Eoc.-Miss.), 24(2) : 37, 6, 35-37; *lobatulus* (Eoc.-Ga.), 21(3) : 72; (Eoc.-Miss.), 21(4) : 105, 16, 23, 24; (Olig.-Miss.), 22(3) : 102; (Plioc.-Alaska), 23(3) : 71, 16, 33; (Olig.-Miss.), 24(1) : 10; *madrugaensis* (Paleoc.-Cuba), 24(4) : 87, 15, 10-12; *martinezensis* (Eoc.-Peru), 25(4) : 83, 14, 27; *mauricensis* (Eoc.-Ala.), 21(1) : 20; (Eoc.-Miss.), 21(4) : 103; 24(2) : 36, 6, 30, 31; *mexicanus* (Eoc.-Peru), 25(4) : 83, 14, 23; *mirificus* (Paleoc.-Cuba), 24(4) : 88, 16, 1-3; *mississippiensis* (Eoc.-Ga.), 21(1) : 10; 21(3) : 72; (Olig.-Miss.), 22(3) : 102; 24(1) : 10; *ouachitaensis* (Eoc.-Miss.), 21(4) : 104, 16, 20; *parki* (Eoc.-Peru), 25(4) : 84, 14, 25; *pippeni* (Olig.-Miss.), 24(1) : 10, 1, 14; *plano-convexa* (Eoc.-Miss.), 21(4) : 104, 16, 18, 19; (Olig.-Miss.), 22(3) : 102; *praeconcentricus* (Eoc.-Ga.), 21(1) : 10, 2, 14; *praecursorius* (Paleoc.-Ark.), 22(2) : 65, 11, 20, 21; (Paleoc.-Cuba), 24(4) : 89; *pseudoungerianus* (Eoc.-Ga.), 21(3) : 73; *refulgens* (Plioc.-Alaska), 23(3) : 71; *reprimatus* (Paleoc.-Cuba), 24(4) : 87, 15,

- 13, 14; **RHSmithi**, 25(4) : 101; *sassei* (Eoc.-Ala.) 21(1) : 20; (Eoc.-Miss.), 24(2) : 36, 6, 34; *stephensoni* (Cret.-Trinidad), 23(2) : 51, 12, 15; *umbilicatus*, 25(4) : 101; *venezuelanus* (Eoc.-Peru), 25(4) : 83, 14, 19; *vicksburgensis* (Olig.-Miss.), 22(3) : 102, 16, 30-32; *vulgaris* (Paleoc.-Ark.), 22(2) : 65, 11, 22, 23; (Paleoc.-Cuba), 24(4) : 89; *westi* (Eoc.-Ga.), 21(3) : 72, 11, 15; (Eoc.-Miss.), 24(2) : 36, 6, 32, 33; *yazooensis* (Eoc.-Miss.), 21(4) : 104, 16, 16, 17; (Olig.-Miss.), 22(3) : 102.
- Clavulinoides aspera* (Cret.-Venezuela), 23(1) : 7, 2, 15; (Cret.-Trinidad), 23(2) : 38; var. *whitei* (Cret.-Venezuela), 23(1) : 7, 2, 14; *eucarinatus* (Eoc.-Peru), 25(4) : 77, 13, 21; *havanensis* (Eoc.-Peru), 25(4) : 77, 13, 20; *midwayensis* (Paleoc.-Ark.), 22(2) : 46, 7, 3-6; *trilatera* (Cret.-Venezuela), 23(1) : 6, 2, 13.
- Coleites* and Its Species, The Genus, 24(4) : 81.
- Coleites*, 24(4) : 81; **abuillotensis** (Eoc.-Haiti), 24(4) : 83, 14, 1, 2; *cancellatus* (Eoc.-Cuba; Paleoc.-Guatemala), 24(4) : 83, 13, 10, 11; *danicus* (Paleoc.-Sweden), 24(4) : 82, 13, 9; **guatemalensis** (Eoc.-Guatemala), 24(4) : 84, 14, 3-5; *laevigatus* (Eoc.-Ala.), 24(4) : 82, 13, 5-8; **pasionensis** (Eoc.-Guatemala), 24(4) : 84, 14, 6; *reticulosus* (Paleoc.-Cuba), 24(4) : 81, 13, 1-4; 89.
- Colomia*, A New Genus from the Upper Cretaceous of Cuba, 24(1) : 12.
- Colomia**, 24(1) : 12; **cretacea** (Cret.-Cuba), 24(1) : 12, 2, 13-15.
- Conicospirillina atlantica* (Rec.-Atl.), 23(4) : 91, 20, 8.
- Cornuspira byramensis* (Olig.-Miss.), 22(3) : 84, 14, 22; 24(1) : 8; *olygogyra* (Eoc.-Miss.), 21(4) : 84, 13, 23, 24; *planorbis* (Plioc.-Alaska), 23(3) : 62.
- [Cretaceous]. A New *Pseudoclavulina* from the Upper Cretaceous of Alabama, 21(3) : 74. Some Foraminifera of Woodbine Age from Texas, Mississippi, Alabama and Georgia, 22(3) : 71. A Foraminiferal Fauna from the Santa Anita Formation of Venezuela, 23(1) : 1. Two New Species of Lower Cretaceous Foraminifera from Florida, 23(2) : 29. Further Notes on the Cretaceous Foraminifera of Trinidad, 23(2) : 31. Some New Foraminifera from the American Cretaceous, 23(3) : 53. Two New Species of Foraminifera from the Arkadelphia Marl of Arkansas, 23(3) : 56. A New Name for *Proteonina testacea* from the Lower Cretaceous of the Netherlands, 23(3) : 72. *Colomia*, A New Genus from the Upper Cretaceous of Cuba, 24(1) : 12. *Globigerinelloides*, A New Genus of the Globiger-

- inidae, 24(2) : 42. A Foraminiferal Fauna from the New Almaden District, California, 24(4) : 90.
- Criboelphidium**, 24(1) : 18; *kugleri* (Rec.-Trinidad), 24(1) : 18, 4, 4; *limosum* (Rec.-Trinidad), 24(1) : 19, 4, 7; *salsum* (Rec.-Trinidad), 24(1) : 19, 4, 6; *trinitatensis* (Rec.-Trinidad), 24(1) : 20, 4, 8; *vadescens* (Rec.-Trinidad), 24(1) : 18, 4, 5.
- Cribropyrgo*, and a New Species of *Rotalia*, A New Genus, 22(4) : 119.
- Cribropyrgo**, 22(4) : 119; *robusta* (Rec.-Cuba.), 22(4) : 119, 20, 7-9.
- Crirostomoides trinitatensis* (Cret.-Venezuela), 23(1) : 4, 1, 6, 7; (Cret.-Trinidad), 23(2) : 37.
- Cristellaria americana*, var. *grandis*, 22(4) : 128; *globosa*, 22(4) : 131; *globulosa*, 22(4) : 131; *incrassata*, 22(4) : 127; *josephina*, var. *umbonata*, 25(4) : 102; *limbata*, 25(4) : 102; *rotulata*, var. *umbonata*, 25(4) : 102; *translucida*, 25(4) : 102.
- Cristellaria (Robulus) grandis*, 22(4) : 128.
- [Cuba]. The Species of *Globigerina* Described by d'Orbigny in 1839 from Shore Sands of Cuba, 21(3) : 74. Some Cuban Foraminifera of the Genus *Rotalia*, 23(2) : 23. *Colomia*, A New Genus from the Upper Cretaceous of Cuba, 24(1) : 12. Some Paleocene Foraminifera from the Madruga Formation of Cuba, 24(3) : 68. Additional Species of Paleocene Foraminifera from the Madruga Formation of Cuba, 24(4) : 85. Some Cuban Species of *Globorotalia*, 25(2) : 26.
- Cuneolina walteri* (Cret.-Fla.), 23(2) : 30, 10, 4, 5.
- Cyclammina deformis* (Eoc.-Peru), 25(4) : 76, 13, 15; *elegans* (Cret.-Venezuela), 23(1) : 4, 1, 8; *garcilassoi* (Cret.-Venezuela), 23(1) : 5, 1, 13-17; *schencki* (Cret.-Venezuela), 23(1) : 5, 1, 18.
- Dentalina aculeata* (Paleoc.-Ark.), 22(2) : 51, 8, 9; *californica* (Plioc.-Calif.), 22(2) : 66, 12, 3-5; *catenula* (Cret.-Trinidad), 23(2) : 41, 11, 8; *colei* (Paleoc.-Ark.), 22(2) : 49, 8, 2; *communis* (Eoc.-Ga.), 21(3) : 59, 9, 17; (Mioc.-Ecuador), 24(3) : 53, 9, 10; *confluens* (Cret.-Venezuela), 23(1) : 11, 3, 14, 15; *consobrina* (Cret.-Venezuela), 23(1) : 11, 3, 16, 17; *eocenica* (Paleoc.-Ark.), 22(2) : 50, 8, 4, 5; *gardnerae* (Paleoc.-Ark.), 22(2) : 49, 8, 1; (Paleoc.-Cuba), 24(3) : 69, 11, 3; *gigantea* (Jur.-Poland), 25(1) : 1; *gracilis* (Cret.-Trinidad), 23(2) : 41, 11, 9; *hantkeni* (Eoc.-Ga.), 21(3) : 59, 9, 16; *inepta* (Paleoc.-Ark.), 23(4) : 83,

18, 5; *insulsa* (Paleoc.-Ark.), 23(4) : 84, 18, 6, 7; *jacksonensis* (Eoc.-Ga.), 21(3) : 58, 9, 15; *jamaicensis* (Mioc.-Jamaica), 22(4) : 120; *jarvisi* (Mioc.-Jamaica), 22(4) : 120; *mauricensis* (Eoc.-Miss.), 21(4) : 86, 14, 3; *megalopolitana* (Cret.-Venezuela), 23(1) : 11, 3, 11-13; (Cret.-Trinidad), 23(2) : 41; *mucronata* (Paleoc.-Ark.), 22(2) : 51, 8, 12; (Eoc. & Olig.-Miss.), 24(1) : 11; *naheolensis* (Paleoc.-Ark.), 22(2) : 50, 8, 8; *nasuta* (Paleoc.-Ark.), 22(2) : 50, 8, 6, 7; *plummerae* (Paleoc.-Ark.), 22(2) : 50, 8, 3; *praecatesbyi* (Olig.-Miss.), 24(1) : 8; *pseudo-involvens* (Olig.-Miss.), 22(3) : 85, 14, 24, 25; **pseudo-nasuta** (Paleoc.-Ark.), 22(2) : 51, 8, 13, 14; *pseudo-obliquestriata* (Paleoc.-Ark.), 22(2) : 50, 8, 18; *recta*, 22(4) : 125, 126; *soluta* (Olig.-Miss.), 22(3) : 85, 14, 26; sp. (Eoc.-Ala.), 21(1) : 13, 3, 13, 14; sp. A (Eoc.-Miss.), 21(4) : 86, 14, 4; sp. B (Eoc.-Miss.), 21(4) : 87, 14, 5; sp. A (Paleoc.-Ark.), 22(2) : 51, 8, 10; sp. B (Paleoc.-Ark.), 22(2) : 51, 8, 15, 16; sp. C (Paleoc.-Ark.), 22(2) : 52, 8, 17; sp. (Olig.-Miss.), 22(3) : 85, 14, 27; sp. A (Eoc.-Miss.), 24(2) : 29, 5, 18; sp. B (Eoc.-Miss.), 24(2) : 29, 5, 19; sp. A (Mioc.-Ecuador), 24(3) : 53, 9, 11; sp. B (Mioc.-Ecuador), 24(3) : 53, 9, 12; sp. (Jur.-Poland), 25(1) : 10, 3, 3; sp. (Eoc.-Peru), 25(4) : 79, 14, 4.

Dentalina (*Nodosaria*) *pyriformis*, 25(4) : 102.

[Descriptive]. A New Species of *Globobulimina* from the Western Atlantic, 21(3) : 73. A New *Pseudoclavulina* from the Upper Cretaceous of Alabama, 21(3) : 74. Some New Species and Varieties of Foraminifera from the Pliocene of Timms Point, California, 22(2) : 65. Some Foraminifera of Woodbine Age from Texas, Mississippi, Alabama, and Georgia, 22(3) : 71. A New Species of *Amphistegina* from the Eocene of Ecuador, 22(4) : 117. A New Genus, *Cribrropyrgo*, and a New Species of *Rotalia*, 22(4) : 119. Some Cuban Foraminifera of the Genus *Rotalia*, 23(2) : 23. Two New Species of Lower Cretaceous Foraminifera from Florida, 23(2) : 29. Some New Foraminifera from the American Cretaceous, 23(3) : 53. Two New Species of Foraminifera from the Arkadelphia Marl of Arkansas, 23(3) : 56. A New Species of *Bulimina* from the Eocene, Chira Shale, of Peru, 23(3) : 57. Some New Species of Foraminifera from the Lower Pliocene of the Netherlands, 23(3) : 57. A New Genus and Some New Species of Foraminifera from the Upper Eocene of Ecuador, 23(4) : 77. Some New Foraminifera from the Paleocene of the Southern United States, 23(4) : 81.

New Species and Varieties of Foraminifera from off the South-eastern Coast of the United States, 23(4): 86. Foraminifera from the Red Bluff-Yazoo Section at Red Bluff, Mississippi, 24(1): 1. *Colomia*, A New Genus from the Upper Cretaceous of Cuba, 24(1): 12. Some New Genera and Species of Foraminifera from Brackish Water of Trinidad, 24(1): 15. A New Species of *Nonion* (Foraminifera) from the Woods Hole Region, 24(1): 21. Additional New Species of Arenaceous Foraminifera from Shallow Waters of Trinidad, 24(2): 37. *Globigerinelloides*, A New Genus of the Globigerinidae, 24(2): 42. Additional New Foraminifera from the American Paleocene, 24(2): 43. *Pseudoparrella*, A New Generic Name, and A New Species of *Parrella*, 24(3): 49. Some Paleocene Foraminifera from the Madrugá Formation of Cuba, 24(3): 68. Additional Species of Paleocene Foraminifera from the Madrugá Formation of Cuba, 24(4): 85. Amendment to the Diagnosis of *Lockhartia huntii* Ovey, 24(4): 99. Some Cuban Species of *Globorotalia*, 25(2): 26. *Pavoninoides*, A New Genus of the Miliolidae from Panama, 25(3): 58.

Dimorphina inflexa (Jur.-Poland), 25(1): 1; *tyraica* (Jur.-Poland), 25(1): 1.

Discocyclina (*Asterocyclina*) sp. (Eoc.-Ecuador), 22(4): 118.

Discocyclina (*Discocyclina*) *anconensis* (Eoc.-Ecuador), 22(4): 118, *meroensis* (Eoc.-Ecuador), 22(4): 118; *sheppardi* (Eoc.-Ecuador), 22(4): 118.

Discospirina italica, 21(2): 49; *liburnica*, 21(2): 49; var. *trilobata*, 21(2): 49.

Discorbis alveata (Eoc.-Miss.), 21(4): 100, 16, 1, 2; *arcuato-costata* (Olig.-Miss.), 22(3): 96, 16, 15; 24(1): 9; *assulata* (Eoc.-Ga.), 21(3): 68, 11, 2; (Eoc.-Miss.), 21(4): 100, 15, 34, 35; (Olig.-Miss.), 22(3): 97; *clementiana*, var. *laevigata*, 22(4): 126; var. *rugosa*, 22(4): 126; *cocoaensis* (Eoc.-Miss.), 24(1): 10, 2, 10; *confusa* (Eoc.-Holland), 22(4): 123; *dreheri* (Jur.-Poland), 25(1): 11, 3, 6, 7; *georgiana* (Eoc.-Ga.), 21(3): 68, 11, 5; *globulo-spinosa* (Eoc.-Miss.), 21(4): 100, 15, 32, 33; *hemisphaerica* (Eoc.-Ga.), 21(3): 67, 11, 3; (Eoc.-Miss.), 21(4): 100, 15, 30, 31; 24(2): 34, 6, 16, 17; *mauricensis* (Eoc.-Miss.), 24(2): 34, 6, 18; *oligocenica* (Olig.-Miss.), 22(3): 97, 16, 9, 14; *orbicularis* (Olig.-Miss.), 22(3): 97; *patellinaformis* (Eoc.-Holland), 22(4): 123; *petalifera* (Eoc.-Ala.), 21(1): 18, 4, 11; *plana* (Eoc.-Holland), 22(4): 121; *pulchra* (Rec.-Atl.),

23(4): 91, 20, 9; *rutteni* (Eoc.-Holland), 22(4): 121; *subglobosa* (Olig.-Miss.), 22(3): 96, 16, 8; 24(1): 9; *valvulata* (Plioc.-Alaska), 23(3): 67, 16, 8; *yeguaensis* (Eoc.-Ga.), 21(3): 68, 11, 4.

Discorinopsis vadescens (Rec.-Trinidad), 24(1): 20, 4, 9, 10.

Dorothia almadenensis (Cret.-Calif.), 24(4): 94, 16, 11, 12.

Dyocibicides biserialis (Plioc.-Alaska), 23(3): 72, 16, 34, 35.

[Ecuador]. A New Species of *Amphistegina* from the Eocene of Ecuador, 22(4): 117. A New Genus and Some New Species of Foraminifera from the Upper Eocene of Ecuador, 23(4): 77.

A Miocene Foraminiferal Fauna from Ecuador, 24(3): 50.

eggerella, 22(4): 131; *trochoides* (Cret.-Europe, Amer.), 22(1): 18; (Cret.-Trinidad), 23(2): 39.

eggerellina gibbosa, var. *conica*, 22(4): 126; var. *globulosa*, 22(4): 126.

Ellipsoglandulina exponens (Cret.-Venezuela), 23(1): 15, 4, 11; (Cret.-Trinidad), 23(2): 48.

Ellipsonodosaria alexanderi, var. *impensia* (Cret.-Venezuela), 23(1): 15, 4, 21, 22; (Cret.-Trinidad), 23(2): 48, 12, 9, 10; *ewaldi* (Eoc.-Ga.), 21(3): 67; *granti* (Eoc.-Miss.), 24(1): 10; *midwayensis* (Paleoc.-Ark.), 22(2): 61, 10, 25; *paleocenica* (Paleoc.-Ark.), 22(2): 61, 10, 26; *pilulata* (Eoc.-Miss.), 24(1): 5, 2, 8; 10, 2, 8; *plummerae* (Paleoc.-Ark.), 22(2): 61, 10, 24; *pseudoscripta* (Cret.-Trinidad), 23(2): 48; *sagrinensis* (Eoc.-Peru), 25(4): 81, 14, 5; *subnodosa* (Cret.-Trinidad), 23(2): 47; sp. (Eoc.-Ga.), 21(1): 9, 2, 8; sp. (Eoc.-Peru), 25(3): 55, 10, 9.

Ellipsopleurostomella curta (Cret.-Trinidad), 23(2): 47.

Elphidiella arctica (Plioc.-Alaska), 23(3): 65, 15, 20.

Elphidioides, 21(1): 7; *americanus* (Eoc.-Ga.), 21(1): 7, 2, 4.

Elphidium alaskense (Plioc.-Alaska), 23(3): 65, 15, 18, 19; *bartletti* (Plioc.-Alaska), 23(3): 64; *fastigiatum* (Eoc.-Ga.), 21(1): 6, 2, 3; *texanum* (Eoc.-Ga.), 21(3): 63, 10, 13; *twiggsanum* (Eoc.-Ga.), 21(1): 6, 2, 2.

Entosolenia catenulata (Plioc.-Alaska), 23(3): 66, 15, 28; *compressa* (Plioc.-Alaska), 23(3): 66, 15, 25; *globosa* (Eoc.-Ga.), 21(3): 64; *howei* (Eoc.-La. & Miss.), 21(4): 95, 15, 29; (Olig.-Miss.), 22(3): 92; *laevigata* (Eoc.-Ga.), 21(3): 64, 10, 17; (Eoc.-Miss.), 21(4): 95, 15, 28; (Olig.-Miss.), 24(1): 9; *lucida* (Plioc.-Alaska), 23(3): 65, 15, 22; *marginata* (Cret.-Trinidad), 23(2): 45; (Eoc.-Miss.), 24(2): 33; *marginata*, var. (Plioc.-

Alaska), 23(3): 65, 15, 23, 24; var. *lisbonensis* (Eoc.-Miss.), 24(2): 33, 6, 13; *marginato-perforata* (Olig.-Miss.), 22(3): 92, 15, 27; *mauricensis* (Eoc.-Miss.), 24(2): 33, 6, 12; *orbignyana* (Cret.-Trinidad), 23(2): 46; *orbignyana*, var. (Plioc.-Alaska), 23(3): 66, 15, 26; var. *flintii* (Olig.-Miss.), 24(1): 9, 1, 8; var. *perfecto-costata* (Cret.-Trinidad), 23(2): 46, 12, 5; *squamosa* (Plioc.-Alaska), 23(3): 66, 15, 27; (Olig.-Miss.), 24(1): 9.

[Eocene]. A Foraminiferal Fauna from the Twiggs Clay of Georgia, 21(1): 1. A Foraminiferal Fauna from the Lisbon Formation of Alabama, 21(1): 11. The Foraminifera of the Type Locality of the McBean Formation, 21(3): 55. Foraminifera of the Type Locality of the Moodys Marl Member of the Jackson Formation of Mississippi, 21(4): 79. A New Species of *Amphistegina* from the Eocene of Ecuador, 22(4): 117. Some Homonyms in "Foraminifera from the Middle Eocene in the Southern Part of the Netherlands Province of Limburg," 22(4): 120. A New Species of *Bulimina* from the Eocene, Chira Shale, of Peru, 23(3): 57. A new Genus and Some New Species of Foraminifera from the Upper Eocene of Ecuador, 23(4): 77. Foraminifera from the Red Bluff-Yazoo Section at Red Bluff, Mississippi, 24(1): 1. A Foraminiferal Fauna from the Cook Mountain Formation of Mississippi, 24(2): 27. *Anomalina tenuissima* (Reuss), var. *evoluta* van Bellen, A Homonym, 24(3): 76. Foraminifera from the Eocene, Chacra Formation of Peru, 25(3): 49. Foraminifera from the Eocene, Verdun Formation, of Peru, 25(4): 73.

Eouvigerina excavata (Paleoc.-Ark.), 22(2): 58, 10, 16; (Paleoc.-Cuba), 24(4): 89.

Epistomina eocenica (Eoc.-Miss.), 24(1): 11, 2, 9; (Eoc.-Peru), 25(4): 82, 14, 17.

Eponides abatissae, var. *multicameratus*, 25(4): 100; *adventus* (Olig.-Miss.), 24(1): 10, 1, 11; *alabamensis* (Olig.-Miss.), 22(3): 99, 16, 22; *bollii* (Cret.-Venezuela), 23(1): 16, 4, 25; (Cret.-Trinidad), 23(2): 49; *bronnimanni* (Cret.-Venezuela), 23(1): 16; Cret.-Trinidad), 23(2): 49; var. *mera* (Cret.-Venezuela), 23(1): 16, 4, 26; (Cret.-Trinidad), 23(2): 49; *budensis* (Eoc.-Holland), 22(4): 121; *byramensis* (Olig.-Miss.), 22(3): 99, 16, 17, 18; *choctawensis* (Olig.-Miss.), 22(3): 99; *cocoaensis* (Eoc.-Ga.), 21(3): 69; *columbiensis* (Plioc.-Alaska), 23(3): 68, 16, 14, 15; *ecuadorana* (Mioc.-Ecuador), 24(3):

64, 10, 16; *frigidus* (Plioc.-Alaska), 23(3): 67, 16, 10-13; *gratiosus* (Paleoc.-Cuba), 24(3): 71, 11, 14; *guayabalensis*, var. *yeguaensis* (Eoc.-Ala.), 21(1): 18, 4, 14; *haidingerii* (Paleoc.-Ark.), 22(2): 62, 11, 5, 6; *inornata*, 25(4): 101; *jacksonensis* (Eoc.-Ga.), 21(1): 10; (Eoc.-Miss.), 21(4): 101, 16, 3; *mexicanus* (Eoc.-Ala.), 21(1): 18, 4, 12, 13; *minimus* (Eoc.-Ga.), 21(1): 9, 2, 12; (Eoc.-Peru), 25(3): 56, 10, 13; *minutissimus* (Olig.-Miss.), 24(1): 5, 1, 13; 10, 1, 13; *multicameratus*, 25(4): 100; *ouachitaensis* (Olig.-Miss.), 24(1): 10, 1, 12; *plummerae* (Paleoc.-Tex.), 24(2): 44, 8, 9; *simplex*, 25(4): 101; *umbonatus* (Plioc.-Alaska), 23(3): 67; (Eoc.-Peru), 25(4): 81, 14, 13; *vanbelleni* (Paleoc.-Cuba), 24(3): 71, 12, 1-3; *zinkstoki* (Eoc.-Holland, 22(4): 121; sp. (Cret.-Calif.), 24(4): 95, 16, 15.

[Faunal]. A Foraminiferal Fauna from the Twiggs Clay of Georgia, 21(1): 1. A Foraminiferal Fauna from the Lisbon Formation of Alabama, 21(1): 11. The Foraminifera of the Type Locality of the McBean Formation, 21(3): 55. Foraminifera of the Type Locality of the Moodys Marl Member of the Jackson Formation of Mississippi, 21(4): 79. A Foraminiferal Fauna from the Paleocene of Arkansas, 22(2): 45. A Foraminiferal Fauna from the Byram Marl at Its Type Locality, 22(3): 76. A Foraminiferal Fauna from the Santa Anita Formation of Venezuela, 23(1): 1. Further Notes on the Cretaceous Foraminifera of Trinidad, 23(2): 31. A Foraminiferal Fauna from Amchitka Island, Alaska, 23(3): 60. Foraminifera from the Red Bluff-Yazoo Section at Red Bluff, Mississippi, 24(1): 1. A Foraminiferal Fauna from the Cook Mountain Formation of Mississippi, 24(2): 27. A Miocene Foraminiferal Fauna from Ecuador, 24(3): 50. A Foraminiferal Fauna from the New Almaden District, California, 24(4): 90. Upper Jurassic Foraminifera from the Nizniow Limestone of Podole, Poland, 25(1): 1. Foraminifera from the Eocene, Chacra Formation, of Peru, 25(3): 49. Foraminifera from the Eocene, Verdun Formation, of Peru, 25(4): 73.

[Florida]. Two New Species of Lower Cretaceous Foraminifera from Florida, 23(2): 29.

Foraminifera from the Eocene, Chacra Formation, of Peru, 25(3): 49.

Foraminifera from the Eocene, Verdun Formation, of Peru, 25(4): 73.

Foraminifera from the Red Bluff-Yazoo Section at Red Bluff, Mississippi, 24(1) : 1.

Foraminifera of the Type Locality of the McBean Formation, The, 21(3) : 55.

Foraminifera of the Type Locality of the Moodys Marl Member of the Jackson Formation of Mississippi, 21(4) : 79.

Foraminiferal Fauna from Amchitka Island, Alaska, A, 23(3) : 60.

Foraminiferal Fauna from the Byram Marl at Its Type Locality, A, 22(3) : 76.

Foraminiferal Fauna from the Cook Mountain Formation of Mississippi, A, 24(2) : 27.

Foraminiferal Fauna from the Lisbon Formation of Alabama, A, 21(1) : 11.

Foraminiferal Fauna from the New Almaden District, California, A, 24(4) : 90.

Foraminiferal Fauna from the Paleocene of Arkansas, A, 22(2) : 45.

Foraminiferal Fauna from the Santa Anita Formation of Venezuela, A, 23(1) : 1.

Foraminiferal Fauna from the Twiggs Clay of Georgia, A, 21(1) : 1.

Fronicularia alata, var. *sagittula*, 22(4) : 126; *bicornis*, var. *rhomboidalis*, 22(4) : 126; *capitana* (Eoc.-Peru), 25(3) : 51, 9, 9-11; *laevigata*, 22(4) : 126; *linguiformis* (Cret.-Trinidad), 23(2) : 43; *mucronata* (Cret.-Venezuela), 23(1) : 13, 4, 3; (Cret.-Trinidad), 23(2) : 43; *naheolensis* (Paleoc.-Ark.), 22(2) : 55, 9, 23; (Paleoc.-Cuba), 24(4) : 89; *rhomboidalis*, 22(4) : 126; *sagittula*, 22(4) : 126; *sepiolaris*, forma nova *typica*, 22(4) : 126; var. *laevigata*, 22(4) : 126; *tenuissima* (Eoc.-Miss.), 24(1) : 10; *typica*, 22(4) : 126; sp. A (Paleoc.-Ark.), 22(2) : 55, 9, 24; sp. (Cret.-Venezuela), 23(1) : 13, 4, 4.

Further Notes on the Cretaceous Foraminifera of Trinidad, 23(2) : 31.

Gaudryina aequa (Rec.-Atl.), 23(4) : 87, 18, 18-21; *almadenensis* (Cret.-Calif.), 24(4) : 92, 16, 5; *althi* (Jur.-Poland), 25(1) : 7, 2, 7; *bentonensis* (Cret.-Trinidad), 23(2) : 38; *bukowiensis* (Jur.-Poland), 25(1) : 8, 2, 10, 11; *crassa*, 22(4) : 127, *dead-ericki* (Cret.-Ark.), 23(3) : 56, 13, 8-10; *exilis* (Rec.-Trinidad), 24(2) : 40, 7, 15, 16; *foeda* (Cret.-Trinidad), 23(2) : 38; *geo-*

metrica (Eoc.-Ala.), 21(1) : 11; *gracilis*, 22(4) : 127; *gradata*, var. *crassa*, 22(4) : 127; var. *gracilis*, 22(4) : 127; *hedbergi* (Cret.-Venezuela), 23(1) : 6, 2, 7; *jacksonensis* (Eoc.-Ga.), 21(3) : 57; *jurassica* (Jur.-Poland), 25(1) : 8, 2, 8, 9; *laevigata* (Cret.-Venezuela), 23(1) : 6, 2, 3-6; *vadaszi* (Jur.-Poland), 25(1) : 7, 2, 5, 6.

Gaudryina (*Pseudogaudryina*) *jacksonensis* (Eoc.-Miss.), 24(1) : 10; *pyramidata* (Cret.-Trinidad), 23(2) : 38.

Gaudryina (*Siphogaudryina*) *glabrata* (Olig.-Miss.) 22(3) : 78, 13, 16, 17; *youngi* (Olig.-Miss.), 22(3) : 77, 13, 15; 24(1) : 5.

Genus *Ceratobulimina* and Its Species, The, 22(4) : 107.

Genus *Coleites* and Its Species, The, 24(4) : 81.

Genus *Hauerina* and Its Species, The, 22(1) : 2.

Genus *Sigmoilina* and Its Species, The, 22(2) : 29.

Genus *Sphaeroidina* and Its Species, The, 25(1) : 11.

[Georgia]. A Foraminiferal Fauna from the Twiggs Clay of Georgia, 21(1) : 1. The Foraminifera of the Type Locality of the McBean Formation, 21(3) : 55. Some Foraminifera of Woodbine Age from Texas, Mississippi, Alabama, and Georgia, 22(3) : 71.

Glandulina ovata (Eoc.-Miss.), 21(4) : 89, 14, 7; 24(2) : 31, 5, 29.

Globigerina Described by d'Orbigny in 1839 from Shore Sands of Cuba, The Species of, 21(3) : 74.

Globigerina Described between 1839 and 1850, The Species of, 22(1) : 15.

Globigerina almadenensis (Cret.-Calif.), 24(4) : 95, 16, 18, 19; *bilobata* (Mioc.-Europe), 22(1) : 19, 3, 9; 4, 7-10; *canariensis* Rec.-Atl.), 22(1) : 15, 3, 1; *concinna* (Mioc.-Europe), 22(1) : 20, 3, 10-12; 4, 11-15; *confluens* (Cret.-Germany), 22(1) : 18; *cretacea* (Cret.-Europe, Amer.), 22(1) : 17, 3, 5; 4, 5, 6; *danvillensis* (Eoc.-Miss.), 21(4) : 102, 16, 11, 12; *dutertrei* (Rec.-Cuba), 21(3) : 75, 12, 8, 11; *elevata* (Cret.-Europe), 22(1) : 18, 3, 4; *foveolata* (Cret.-Africa), 22(1) : 18; *globosa* (Cret.-Germany), 22(1) : 18; *inflata* (Rec.-Atl.), 22(1) : 16, 3, 3; 4, 1-4; (Eoc. & Olig.-Miss.), 24(1) : 11; *pachyderma* (Plioc.-Alaska), 23(3) : 70, 16, 27, 28; *pseudobulloides* (Paleoc.-Ark.), 22(2) : 64; (Paleoc.-Cuba), 24(4) : 89; (Eoc.-Peru), 25(3) : 57, 10, 15; *quadrilatera* (Plioc.-Alaska), 23(3) : 70, 16, 29, 30; *quadrilobata* (Plioc., Mioc.-Europe), 22(1) : 19, 3, 7; *regularis* (Mioc.-Austria), 22(1) : 19, 3, 6; *siphonifera* (Rec.-Cuba), 21(3) : 75, 12, 7, 10; *triloculinooides* (Paleoc.-Ark.), 22(2) : 64;

- trochoides* (Cret.-Europe, Amer.), 22(1) : 18; sp. (Cret.-Calif.), 24(4) : 96, 16, 17.
- Globigerina* (*Globotruncana*) *marginata*, subsp. *linnaeana*, 22(4) : 128.
- Globigerina* (*Rosalina*, *Globotruncana*) *linneana*, 22(4) : 128.
- Globigerinella hirsuta* (Rec.-Atl.), 22(1) : 16, 3, 2.
- Globigerinelloides*, A New Genus of the Globigerinidae, 24(2) : 42.
- Globigerinelloides**, 24(2) : 42; *algeriana* (Cret.-Algeria), 24(2) : 43, 8, 4-6.
- Globigerinoides rubra* (Rec.-Cuba), 21(3) : 75, 12, 6, 9; *triloba* (Mioc. to Rec.), 22(1) : 20, 3, 8; 4, 16-18.
- Globobulimina* from the Western Atlantic, A New Species of, 21(3) : 73.
- Globobulimina caribbea* (Rec.-Cuba), 21(3) : 73, 12, 1-3; *hannai* (Eoc.-Peru), 25(4) : 79, 14, 8.
- Globorotalia*, Some Cuban Species of, 25(2) : 26.
- Globorotalia*, 25(2) : 26; **albeari** (Paleoc.-Cuba), 25(2) : 33, 6, 13-15; **almadenensis** (Cret.-Calif.), 24(4) : 98, 16, 24; **californica** (Cret.-Calif.), 24(4) : 96, 16, 22, 23; **capdevilensis** (Eoc.-Cuba), 25(2) : 32, 6, 10-12; **centralis** (Eoc.-Ga.), 21(3) : 71; **cocoaensis** (Eoc.-Ga.), 21(3) : 71, 11, 13; (Eoc.-Miss.), 21(4) : 102, 16, 13; **compressa** (Paleoc.-Cuba), 25(2) : 34, 6, 19-21; **crassata** (Eoc.-Ga.), 21(3) : 71; var. **aequa** (Paleoc.-Cuba), 24(4) : 89; **decorata** (Cret.-Calif.), 24(4) : 97, 16, 21; **fohsi** (Olig.-Cuba), 25(2) : 30, 5, 14-16; **inconspicua** (Eoc.-Miss.), 24(2) : 36, 6, 26, 27; **lehneri** (Eoc.-Cuba), 25(2) : 32, 6, 7-9; **membranacea** (Cret.-Trinidad), 23(2) : 50; (Paleoc., Cret.-Cuba), 25(2) : 34, 6, 16-18; (Eoc.-Peru), 25(3) : 57, 10, 16; **menardii** (Rec.-Cuba), 25(2) : 29, 5, 4-6; var. **fimbriata** (Rec.-Cuba), 25(2) : 30, 5, 7; var. **multicamerata** (Rec., Mioc.-Cuba), 25(2) : 30, 5, 8-13; **palmerae** (Eoc.-Cuba), 25(2) : 31, 6, 4-6; **praemenardii** (Mioc.-Ecuador), 24(3) : 66, 10, 21, 22; (Olig.-Cuba), 25(2) : 31, 5, 17-19; 6, 1-3; **tumida** (Rec.-Cuba), 25(2) : 28, 5, 1-3; **wilcoxensis** (Eoc.-Miss.), 24(2) : 36, 6, 24, 25; var. **acuta** (Paleoc.-Cuba), 24(4) : 89; sp. (Cret.-Venezuela), 23(1) : 17, 4, 29.
- Globorotalia* (*Truncorotalia*) *aragonensis* (Eoc.-Cuba), 25(2) : 38, 7, 13-15; **brödermanni** (Eoc.-Cuba), 25(2) : 40, 7, 22-24; **crassata** (Eoc.-Cuba), 25(2) : 37, 7, 4-6; var. **aequa** (Paleoc.-Cuba), 25(2) : 37, 7, 7-9; var. **densa** (Eoc.-Cuba), 25(2) : 38, 7, 10-12; **hirsuta** (Plioc.-Cuba), 25(2) : 36, 7, 1-3; **spinulosa** (Eoc.-

- Cuba), 25(2) : 40, 8, 1-3; *truncatulinoides* (Rec.-Cuba), 25(2) : 35, 6, 22-24; *velascoensis* (Paleoc., Cret.-Cuba), 25(2) : 41, 8, 4-6; *wilcoxensis* (Paleoc.-Cuba), 25(2) : 39, 7, 16-18; var. *acuta* (Paleoc.-Cuba), 25(2) : 39, 7, 19-21.
- Globorotalia* (*Turborotalia*) *centralis* (Eoc.-Cuba), 25(2) : 44, 8, 19-21; *cerro-azulensis* (Eoc.-Cuba), 25(2) : 42, 8, 10-12; *mayeri* (Olig.-Cuba), 25(2) : 44, 8, 16-18; *oceanica* (Rec.-Cuba), 25(2) : 43, 8, 13-15; *punctulata* (Rec.-Cuba), 25(2) : 42, 8, 7-9.
- Globotruncana* *arca* (Cret.-Venezuela), 23(1) : 17, 4, 31; (Cret.-Trinidad), 23(2) : 50, 12, 11; *conica* (Cret.-Trinidad), 23(2) : 50, 12, 12; var. *plicata* (Cret.-Trinidad), 23(2) : 50, 12, 13; *globigerinoides*, 22(4) : 128; *lapparenti*, 22(4) : 128, 130; subsp. *lapparenti*, 22(4) : 130; *linnei*, var. *stuarti*, 22(4) : 127; *renzi*, 22(4) : 128.
- Globulina* *alabamensis* (Olig.-Miss.), 22(3) : 87, 15, 4; 24(1) : 8; *gibba* (Eoc.-Ga.), 21(1) : 4, 1, 9; (Eoc.-Ala.), 21(1) : 14, 3, 22; (Eoc.-Ga.), 21(3) : 60; (Eoc.-Miss.), 21(4) : 88, 14, 13; (Paleoc.-Ark.), 22(2) : 56, 10, 3; (Olig.-Miss.), 22(3) : 87; 24(1) : 8; (Eoc.-Miss.), 24(2) : 31, 5, 28; var. *punctata* (Eoc.-Miss.), 21(4) : 88, 14, 14; (Olig.-Miss.), 24(1) : 8; *inaequalis* (Eoc.-Ga.), 21(1) : 4, 1, 10; (Eoc.-Miss.), 21(4) : 89, 14, 10; (Olig.-Miss.), 22(3) : 87; *lacrima*, var. *subsphaerica* (Cret.-Trinidad), 23(2) : 43; *minuta* (Eoc.-Ga.), 21(3) : 60, 10, 2; (Eoc.-Miss.), 21(4) : 89, 14, 16; (Paleoc.-Ark.), 22(2) : 56, 10, 5; *münsteri* (Eoc.-Ga.), 21(1) : 4, 1, 11; 21(3) : 60, 10, 1; *rotundata* (Eoc.-Ala.), 21(1) : 14; (Paleoc.-Ark.), 22(2) : 56, 10, 4; sp. (Eoc.-Ala.), 21(1) : 14, 3, 23.
- Glomospira* *charoides* (Eoc.-Peru), 25(4) : 75, 13, 12; var. *corona* (Cret.-Trinidad), 23(2) : 36; *gordialis*, var. *diffundens* (Cret.-Trinidad), 23(2) : 36.
- Gümbelina* *cubensis* (Eoc.-Ga.), 21(3) : 63, 10, 14; (Olig.-Miss.), 22(3) : 90; (Eoc. & Olig.-Miss.), 24(1) : 11; *excolata* (Cret.-Trinidad), 23(2) : 44, 11, 13, 14; *garretti* (Eoc.-Ala.) 21(1) : 16, 4, 1; *mauriciana* (Eoc.-Ala.), 21(1) : 16, 4, 2; *midwayensis* (Paleoc.-Ark.), 22(2) : 58, 10, 15; *plummerae* (Cret.-Trinidad), 23(2) : 44; *ultimatumida* (Cret.-Venezuela), 23(1) : 13, 4, 36; (Cret.-Trinidad), 23(2) : 44; *venezuelana* (Eoc.-Miss.), 21(4) : 94, 15, 9.
- Gümbelitria* *columbiana* (Eoc.-Ala.), 21(1) : 16, 4, 3; (Eoc.-Ga.), 21(3) : 64.
- Guttulina* *adhaerens* (Cret.-Trinidad), 23(2) : 43; *byramensis*

(Olig.-Miss.) 22(3) : 86, 15, 3; 24(1) : 8; *caudata* (Eoc.-Ga.), 21(1) : 3, 1, 8; (Eoc.-Ala.), 21(1) : 14, 3, 18, 19; (Eoc.-Ga.), 21(3) : 60, 9, 24; *frankei* (Olig.-Miss.), 24(1) : 8, 1, 7; *hantkeni* (Paleoc.-Ark.), 22(2) : 56, 10, 2; (Eoc.-Miss.), 24(2) : 30, 5, 27; *irregularis* (Eoc.-Ala.), 21(1) : 14, 3, 20, 21; (Eoc.-Ga.), 21(3) : 59, 9, 22; (Eoc.-Miss.), 21(4) : 87, 14, 9; 24(2) : 31, 5, 26; *lisbonensis* (Eoc.-Ala.), 21(1) : 13, 3, 16, 17; *problema* (Eoc.-Ga.), 21(3) : 60, 9, 25; (Paleoc.-Ark.), 22(2) : 56, 10, 1; *similis* (Jur.-Poland), 25(1) : 10, 2, 19; *spicaeformis* (Eoc.-Ga.), 21(3) : 60, 9, 23; (Eoc.-Miss.), 21(4) : 88, 14, 12; *trigonalis* (Cret.-Trinidad), 23(2) : 43; *yamazakii* (Plioc.-Alaska), 23(3) : 63, 15, 12; sp. (Plioc.-Alaska), 23(3) : 63, 15, 13.

Gypsina globula (Eoc.-Miss.), 21(4) : 105, 16, 25.

Gyroidina beisseli (Cret.-Venezuela), 23(1) : 16, 4, 27; **byramensis** (Olig.-Miss.), 22(3) : 98, 16, 19-21; 24(1) : 9; *danvillensis* (Eoc.-Miss.), 24(1) : 11; *depressa* (Cret.-Venezuela), 23(1) : 15; (Cret.-Trinidad), 23(2) : 48; var. *colombiana* (Cret.-Venezuela), 23(1) : 15, 4, 24; (Cret.-Trinidad), 23(2) : 48; *girardana* (Cret.-Trinidad), 23(2) : 48; (Eoc.-Peru), 25(4) : 81, 14, 12; var. *perampla* (Eoc.-Peru), 25(4) : 81, 14, 15; *globosa* (Cret.-Venezuela), 23(1) : 15, 4, 23; (Cret.-Trinidad), 23(2) : 48; **madrugensis** (Paleoc.-Cuba), 24(4) : 85, 14, 10-12; *multilocula* (Mioc.-Ecuador), 24(3) : 64, 10, 14, 15; *scita* (Eoc.-Peru), 25(3) : 56, 10, 12; *soldanii* (Mioc.-Ecuador), 24(3) : 64, 10, 13; var. *octocamerata* (Eoc.-Ga.), 21(3) : 69; (Eoc.-Miss.), 21(4) : 101, 16, 4; 24(2) : 34, 6, 20, 21; *subangulata* (Paleoc.-Ark.), 22(2) : 62, 11, 2-4; *vicksburgensis* (Olig.-Miss.), 24(1) : 10, 1, 10; sp. (Cret.-Calif.), 24(4) : 95, 16, 16.

Gyromorphina, 22(4) : 131.

Hantkenina alabamensis (Olig.-Miss.), 24(1) : 10; *longispina* (Eoc.-Ga.), 21(3) : 70, 11, 11, 12.

Haplophragmium podolicum (Jur.-Poland), 25(1) : 1; **salsum** (Rec.-Trinidad), 24(1) : 16, 3, 10-13.

Haplophragmoides canui (Jur.-Poland), 25(1) : 4, 1, 3; *carinatum* (Eoc.-Peru), 25(4) : 76, 13, 13; *columbiense*, var. *evolutum* (Rec.-Pac.), 24(3) : 76; *coronata* (Cret.-Trinidad), 23(2) : 36; *eggeri* (Cret.-Venezuela), 23(1) : 4, 1, 5; *flagleri*, var. *trinitatis* (Cret.-Trinidad), 23(2) : 37; *glabra* (Cret.-Trinidad), 23(2) : 36; *mauricensis* (Eoc.-Peru), 25(3) : 51, 9, 4; sp. (Eoc.-Ala.), 21(1) : 11, 3, 2.

Hastigerinella jarvisi (Eoc.-Peru), 25(4) : 83, 14, 22; sp. (Cret.-Calif.), 24(4) : 96, 16, 20.

Hauerina and Its Species, The Genus, 22(1) : 2.

Hauerina, 22(1) : 2; *alata* (Tert.-Germany), 22(1) : 14; *alternans* (Plioc.-Italy), 22(1) : 14; *antiqua* (Cret.-Germany), 22(1) : 14; *atlantica* (Rec.-Atl.), 22(1) : 8, 1, 25, 26; *borealis* (Rec.-Atl.), 22(1) : 15; *bradyi* (Rec.-Pac.), 22(1) : 11, 2, 14, 20, 21; *byramensis* (Olig.-Miss., Ala.), 22(1) : 3, 2, 15; (Olig.-Miss.), 22(3) : 82, 14, 12; *circinata* (Rec.-Pac.) 22(1) : 15; "*complanata*" (Rec.-Ceylon), 22(1) : 10, 2, 7; *compressa* (Mioc.-Austria), 22(1) : 4, 1, 15; *desertorum* (Eoc.-Libya), 22(1) : 2, 1, 5, 6; *diversa* (Rec.-Pac.), 22(1) : 11, 2, 16-19; *exigua* (Rec.-Pac.), 22(1) : 14; *fragilissima* (Rec.-Indo-Pac.), 22(1) : 9, 2, 1-6, 8; (Olig.-Miss.), 22(3) : 82; *glabrata* (Plioc.-Fla.), 22(1) : 7, 1, 19; *howelli* (Rec.-Cuba, Fla.), 22(1) : 7, 1, 21; 2, 10; *hyalina* (Rec.-Atl.), 22(1) : 14; *imprimata* (Olig.-Miss.), 22(1) : 3, 1, 7, 8; *inconstans* (Rec.-Pac., Atl.), 22(1) : 15; *inflata* (Plioc.-Italy), 22(1) : 7, 1, 20; *intermedia* (Tert.-Australia), 22(1) : 15; *involuta* (Rec.-Pac.), 22(1) : 13, 2, 25-28; *miocenica* (Mioc.-Fla.), 22(1) : 6, 1, 17; 2, 11; *nautiliformis* (Plioc.-Italy), 22(1) : 14; *occidentalis* (Rec.-Atl.), 22(1) : 9, 1, 22-24; *orientalis* (Rec.-Pac.), 22(1) : 12, 2, 22-24; *ornatissima* (Mioc.-Hungary, Austria), 22(1) : 5, 1, 16; *pacifica* (Rec.-Pac.), 22(1) : 11, 2, 12, 13; *parri* (Olig.-Australia), 22(1) : 6, 1, 18; *sansebastianensis* (Olig.-Porto Rico), 22(1) : 4, 1, 12; *simplex* (Eoc.-Somaliland), 22(1) : 3, 2, 9; *speciosa* (Mioc., Rec.), 22(1) : 5, 1, 13, 14; *tateana* (Olig.-Australia), 22(1) : 4, 1, 9, 10; sp. (Olig.-Miss.), 22(1) : 13, 1, 11; sp. (Eoc.-Somaliland), 22(1) : 14.

Heronallenia vicksburgensis (Olig.-Miss.), 24(1) : 9.

[Holland]. Some Homonyms in "Foraminifera from the Middle Eocene in the Southern Part of the Netherlands Province of Limburg," 22(4) : 120. Some New Species of Foraminifera from the Lower Pliocene of the Netherlands, 23(3) : 57. A new Name for *Proteonina testacea* from the Lower Cretaceous of the Netherlands, 23(3) : 72.

Homonyms in "Foraminifera from the Middle Eocene in the Southern Part of the Netherlands Province of Limburg," Some, 22(4) : 120.

Homonyms in Foraminifera Erected During 1946, 23(4) : 92.

Homonyms in Foraminifera Erected During 1948 and 1949, 25(4) : 100.

Hormosina globulifera (Eoc.-Peru), 25(4): 75, 13, 6, 7; var. *trinitatensis* (Cret.-Trinidad), 23(2): 35.

[Jurassic]. Upper Jurassic Foraminifera from the Nizniow Limestone of Podole, Poland, 25(1): 1.

Karrerella advena (Olig.-Miss.), 24(1): 8, 1, 2; *baccata*, var. *alaskensis* (Plioc.-Alaska), 23(3): 61, 14, 10, 11; *chilostoma* (Eoc.-Peru), 25(4): 78, 13, 24; *mauricensis* (Eoc.-Miss.), 21(4): 80, 13, 6; sp. (Plioc.-Alaska), 23(3): 61, 14, 8, 9.

Labrospira, 24(1): 15; *columbiensis*, var. *robusta* (Rec.-Pac.), 24(3): 76; *salsa* (Rec.-Trinidad), 24(1): 16, 3, 5, 6.

Lagena acuticosta (Eoc.-Ga.), 21(3): 59, 9, 19; (Eoc.-Miss.), 21(4): 87, 14, 8; (Plioc.-Alaska), 23(3): 63, 15, 8; (Eoc. & Olig.-Miss.), 24(1): 11; *amphora*, var. *cylindrica*, 22(4): 127; *apiculata* (Paleoc.-Ark.), 22(2): 56, 9, 25; (Cret.-Venezuela), 23(1): 13, 4, 6; var. *ovoidea*, 22(4): 127; *auriculata*, var. *costata*, 22(4): 129; *cineta*, 22(4): 125; *costae*, 22(4): 125; *costata* (Eoc.-Ga.), 21(3): 59, 9, 18; (Olig.-Miss.), 22(3): 86; 22(4): 129; (Eoc. & Olig.-Miss.), 24(1): 11; (Eoc.-Miss.), 24(2): 29, 5, 20; *cylindrica*, 22(4): 127; *elliptica*, 25(4): 101; *fenestris-sima* (Eoc.-Miss.), 24(2): 29, 5, 23; *flexa* (Plioc.-Calif.), 22(2): 68, 12, 18-21; *gutta*, 22(4): 131; *guttata*, 22(4): 131; *guttula*, 22(4): 131; *heron-alleni*, 22(4): 125; *hexagona* (Eoc.-Ga.), 21(1): 3, 1, 7; (Olig.-Miss.), 22(3): 86; *hispida*, var. *ovoidea*, 22(4): 127; *laevis* (Eoc.-Ga.), 21(3): 59, 9, 20; (Paleoc.-Ark.), 22(2): 56, 9, 26; *lisbonensis* (Eoc.-Miss.), 24(2): 30, 5, 22; *mississippiensis* (Eoc.-Miss.), 24(2): 30, 5, 24; *mucronata*, 22(4): 127; *neglecta*, 22(4): 125; *nucelloides*, var. *corrosa*, 22(4): 125; *ollula*, 22(4): 131; *ollulata*, 22(4): 131; *ovum*, var. *mucronata*, 22(4): 127; *parvulipora*, 25(4): 102; *pliocenica* (Plioc.-Calif.), 22(2): 68, 12, 22-25; var. *timmsana* (Plioc.-Calif.), 22(2): 68, 12, 15-17; *sidebottomi*, 22(4): 125; *simplex*, 22(4): 125; *striata*, var. *mucronata*, 22(4): 127; *striato-punctata* (Olig.-Miss.), 22(3): 86; *substriata* (Eoc.-Ala.), 21(1): 13, 3, 15; (Eoc.-Miss.), 24(2): 29, 5, 21; *sulcata*, var. *apiculata* (Plioc.-Calif.), 22(2): 68, 12, 12; var. *laevicostata* (Plioc.-Calif.), 22(2): 68, 12, 13, 14; *tetragona* (Eoc.-Ga.), 21(3): 59, 9, 21; *williamsoni* (Plioc.-Alaska), 23(3): 63, 15, 9; sp. A (Paleoc.-Ark.), 22(2): 56, 9, 27; sp. A (Plioc.-Alaska),

- 23(3) : 63, 15, 10; sp. B (Plioc.-Alaska), 23(3) : 63, 15, 11; sp. (Eoc.-Miss.), 24(2) : 30, 5, 25.
- Lagena* (*Fissurina*) *marginata*, subsp. *costata*, 22(4) : 129.
- Lagunculina* *vadescens* (Rec.-Trinidad), 24(1) : 15, 3, 1, 2.
- Lamarckina* *byramensis* (Olig.-Miss.), 22(3) : 98, 16, 12, 13; 24(1) : 9; *glabrata* (Olig.-Miss.), 22(3) : 97; 24(1) : 9; **paleo-
cenica** (Paleoc.-Tenn.), 24(2) : 44, 8, 8; sp. (Eoc.-Miss.), 21(4) : 100.
- Lasiotrochus*, 22(4) : 131.
- Lenticulina* *frankei*, 22(4) : 127; *incrassata*, 22(4) : 127; *navicula* (Cret.-Venezuela), 23(1) : 10, 3, 7; *polygonalis*, 22(4) : 131; *polygonata*, 22(4) : 131; *theta* (Eoc.-Ala.), 21(1) : 13, 3, 12.
- Lenticulina* (*Nautilus*) *incrassata*, 22(4) : 127.
- Leptodermella* *arenata* (Eoc.-Ga.), 21(1) : 3, 1, 6; **excentrica** (Rec.-Trinidad), 24(2) : 37, 7, 1, 2; *salsa* (Rec.-Trinidad), 24(1) : 15, 3, 3, 4.
- Liebusella* *byramensis* (Olig.-Miss.), 22(3) : 78, 13, 20; var. *turgida* (Olig.-Miss.), 24(1) : 8.
- Lingulina* *naheolensis* Cushman, A New Name, 23(1) : 18.
- Lingulina* **naheolensis** (Paleoc.-Ala.), 23(1) : 18; *wilcoxensis* (Paleoc.-Ark.), 22(2) : 54, 9, 12.
- Lituola* **compressa** (Jur.-Poland), 25(1) : 5, 1, 8-10; **podolica** (Jur.-Poland), 25(1) : 4, 1, 4, 5; **siemiradzkii** (Jur.-Poland), 25(1) : 5, 1, 6, 7.
- Lituotuba* *chirana* (Eoc.-Peru), 25(4) : 76, 13, 10, 11; *incerta* (Jur.-Poland), 25(1) : 3, 1, 2; *lituiformis* (Cret.-Trinidad), 23(2) : 36.
- Lockhartia* *hunti*, 24(4) : 99.
- Loxostomum* *applinae* (Paleoc.-Ark.), 22(2) : 60, 10, 22; (Paleoc.-Cuba), 24(3) : 71, 11, 10; *claibornense* (Eoc.-Ga.), 21(3) : 66, 10, 25; *dalli* (Eoc.-Miss.), 24(1) : 10, 2, 6; **deadericki** (Paleoc.-Ark.), 23(4) : 85, 18, 8-10; var. **exilis** (Paleoc.-Ark.), 23(4) : 85, 18, 11-13; **lammersi** (Plioc.-Holland), 23(3) : 58, 14, 6; *limonense* (Cret.-Trinidad), 23(2) : 46; *plummerae* (Paleoc.-Ark.), 22(2) : 60, 10, 21; *trinitatensis* (Cret.-Trinidad), 23(2) : 46; *vicksburgense* (Olig.-Miss.), 22(3) : 93, 15, 34.
- Marginulina* *abbreviata* (Eoc. & Olig.-Miss.), 24(1) : 11; *bullata* (Cret.-Venezuela), 23(1) : 10, 3, 8; (Cret.-Trinidad), 23(2) : 40; *burgundiae*, subsp. *dentaliniformis*, 22(4) : 127; *cocoaensis* (Eoc.-Ga.), 21(3) : 58, 9, 8-13; (Eoc. & Olig.-Miss.), 24(1) : 11,

- 1, 15; *decorata* (Cret.-Venezuela), 23(1): 11, 3, 5; *dentalini-formis*, 22(4): 127; *distincta* (Paleoc.-Cuba), 24(3): 69, 11, 2; *dubia* (Paleoc.-Ark.), 22(2): 49, 7, 18; *eximia* (Paleoc.-Ark.), 22(2): 48, 7, 17; *fragaria*, var. *texasensis* (Eoc.-Miss.), 21(4): 85, 13, 29; *grata* (Cret.-Venezuela), 23(1): 11, 3, 10; *hamata* (Paleoc.-Ark.), 22(2): 49, 7, 19, 20; *jarvisi* (Cret.-Venezuela), 23(1): 11; (Cret.-Trinidad), 23(2): 41; *karreriana* (Eoc.-Ga.), 21(3): 58, 9, 14; *lalickeri* (Eoc.-Miss.), 24(1): 10; *mexicana*, var. *nudicostata* (Eoc.-Peru), 25(4): 79, 14, 2, 3; *moodyensis* (Eoc.-Miss.), 21(4): 85, 14, 1, 2; *orbignyana*, 22(4): 131; *orbignyi*, 22(4): 131; *radiata* (Jur.-Poland), 25(1): 10, 2, 20, 21; *subrecta* (Paleoc.-Ark.), 22(2): 49, 7, 23; (Eoc.-Miss.), 24(1): 10; *texasensis* (Cret.-Trinidad), 23(2): 41; *trilobata* (Cret.-Trinidad), 23(2): 41; *tuberculata* (Paleoc.-Cuba), 24(3): 69, 11, 5; *villa* (Rec.-Atl.), 23(4): 89, 19, 7, 8; sp. (Paleoc.-Ark.), 22(2): 49, 7, 21, 22; sp. (Cret.-Venezuela), 23(1): 11, 3, 9.
- Marssonella indentata* (Cret.-Venezuela), 23(1): 7, 2, 16; (Cret.-Trinidad), 23(2): 39; *oxycona*, var. *trinitatensis* (Cret.-Venezuela), 23(1): 7, 2, 8; (Cret.-Trinidad), 23(2): 39.
- Martinottiella paleocenica* (Paleoc.-Ala.), 23(4): 82, 18, 1.
[Massachusetts]. A New Species of *Nonion* (Foraminifera) from the Woods Hole Region, 24(1): 21.
- Massilina crusta* (Olig.-Miss.), 22(3): 80, 14, 5; *decorata* (Olig.-Miss.), 24(1): 8; (Eoc.-Peru), 25(4): 78, 13, 29; *jacksonensis* (Eoc.-Miss.), 21(4): 82, 13, 17; var. *punctato-costata* (Eoc.-Miss.), 21(4): 82, 13, 18; *mauricensis* (Eoc.-Miss.), 24(2): 28, 5, 10; *pulchra* (Plioc.-Calif.), 22(2): 65, 12, 1.
- Miliola jacksonensis* (Eoc.-Miss.), 21(4): 83, 13, 13; *saxorum* (Eoc.-Miss.), 21(4): 83, 13, 14.
- Miliolinella robusta* (Olig.-Miss.), 24(1): 2, 1, 3; 8, 1, 3.
- Millettella elongata* (Eoc.-Ga.), 21(1): 2, 1, 5; *eocenica* (Eoc.-Ga.), 21(1): 2, 1, 4.
- Mimosina*, 21(2): 43; *affinis* (Rec.-Pac.), 21(2): 43, 7, 15, 16; *echinata* (Rec.-Pac., Atl., Medit.), 21(2): 44, 7, 20-22; *hystrix* (Rec.-Pac.), 21(2): 44, 7, 17, 18; *pacifica* (Rec.-Pac.), 21(2): 45, 7, 23; *rimosa* (Rec.-Pac.), 21(2): 44, 7, 19.
- [Miocene]. A Miocene Foraminiferal Fauna from Ecuador, 24(3): 50.
- [Mississippi]. Foraminifera of the Type Locality of the Moodys Marl Member of the Jackson Formation of Mississippi, 21(4):

79. Some Foraminifera of Woodbine Age from Texas, Mississippi, Alabama, and Georgia, 22(3): 71. A Foraminiferal Fauna from the Byram Marl at Its Type Locality, 22(3): 76. Foraminifera from the Red Bluff-Yazoo Section at Red Bluff, Mississippi, 24(1): 1. A Foraminiferal Fauna from the Cook Mountain Formation of Mississippi, 24(2): 27.

Mississippina monsouiri (Olig.-Miss.), 22(3): 100, 16, 26.

New Genus and Some New Species of Foraminifera from the Upper Eocene of Ecuador, A, 23(4): 77.

New Genus, *Cribropyrgo*, and a New Species of *Rotalia*, A, 22(4): 119.

New Name for *Proteonina testacea* from the Lower Cretaceous of the Netherlands, A, 23(3): 72.

New Names for Four Homonym Species Described in "Foraminifera in the Gullmar Fjord and the Skagerak," 24(2): 45.

New *Pseudoclavulina* from the Upper Cretaceous of Alabama, A, 21(3): 74.

New Species and Varieties of Foraminifera from off the Southeastern Coast of the United States, 23(4): 86.

New Species of *Amphistegina* from the Eocene of Ecuador, A, 22(4): 117.

New Species of *Bulimina* from the Eocene, Chira Shale, of Peru, A, 23(3): 57.

New Species of *Globobulimina* from the Western Atlantic, A, 21(3): 73.

New Species of *Nonion* (Foraminifera) from the Woods Hole Region, A, 24(1): 21.

Nodellum velascoense (Cret.-Trinidad), 23(2): 35.

Nodobaculariella scita (Olig.-Miss.), 24(1): 2, 1, 5; 8, 1, 5; sp. (Eoc.-Miss.), 21(4): 84, 13, 21, 22.

Nodogenerina advena (Mioc.-Ecuador), 24(3): 55, 10, 3.

Nodosarella coalingensis (Cret.-Trinidad), 23(2): 47; **paleocenica** (Paleoc.-Ark.), 22(2): 60, 10, 23.

Nodosaria affinis (Paleoc.-Ark.), 22(2): 52, 8, 21-24; (Cret.-Venezuela), 23(1): 12, 3, 25; (Cret.-Trinidad), 23(2): 42; (Paleoc.-Cuba), 24(4): 89; *amphioxys* (Paleoc.-Ark.), 22(2): 52, 8, 25; *annulifera*, 22(4): 127, 128; *bradyi*, 25(4): 102; *caudata*, 22(4): 128; *ewaldi* (Eoc.-Miss.), 21(4): 87, 14, 6; *gracilitatis* (Cret.-Trinidad), 23(2): 42, 11, 10, 11; *laevigata*, 22(4): 128; *latejugata* (Paleoc.-Ark.), 22(2): 52, 8, 19, 20;

limbata (Cret.-Trinidad), 23(2): 41; *limonensis* (Cret.-Venezuela), 23(1): 12, 3, 26; *longiscata* (Paleoc.-Ark.), 22(2): 52, 8, 26; (Paleoc.-Cuba), 24(4): 89; *marcki* (Cret.-Trinidad), 23(2): 41; *monile* (Cret.-Venezuela), 23(1): 12, 3, 18-21; *paupercula* (Cret.-Venezuela), 23(1): 12, 3, 22-24; (Cret.-Trinidad), 23(2): 41; *praecatesbyi* (Olig.-Miss.), 22(3): 85, 14, 28, 29; *pyriformis*, 25(4): 102; *rugosa*, 23(4): 93; *tryloniformis*, 25(4): 102.

Nodosaria (*Dentalina*) *bradyi*, 25(4): 102.

Nodosaria (*Glandulina*) *laevigata*, 22(4): 128.

Nodosaria (*Vaginulina*) *caudata*, 22(4): 128.

Nonion (Foraminifera) from the Woods Hole Region, A New Species of, 24(1): 21.

Nonion advenum (Eoc.-Ga.), 21(1): 5, 1, 15; 21(3): 61, 10, 9; (Eoc.-Miss.), 21(4): 91, 15, 1; (Olig.-Miss.), 22(3): 88; *affine* (Olig.-Miss.), 22(3): 88, 15, 8; *agrestum* (Mioc.-Ecuador), 24(3): 53, 9, 14; *alabamense* (Olig.-Miss.), 22(3): 88, 15, 9, 10; *costiferum* (Mioc.-Ecuador), 24(3): 54; *danvillense* (Eoc.-Miss.), 21(4): 92, 15, 4; (Eoc. & Olig.-Miss.), 24(1): 11; (Eoc.-Miss.), 24(2): 32, 6, 5; *decoratum* (Olig.-Miss.), 22(3): 88; *florinense* (Eoc.-Ala.), 21(1): 15, 3, 30, 31; (Eoc.-Miss.), 24(2): 32, 6, 2; *inexcavatum* (Eoc.-Ga.), 21(1): 5, 1, 16; 21(3): 62, 10, 10; (Eoc.-Miss.), 21(4): 92, 15, 3; *mauricensis* (Eoc.-Ala.), 21(1): 15, 3, 27, 28; (Eoc.-Miss.), 24(2): 32, 6, 3; *micrum* (Eoc.-Ga.), 21(3): 62, 10, 11; *obducum* (Mioc.-Ecuador), 24(3): 53, 9, 13; *ornatum* (Eoc.-Holland), 22(4): 121; *planatum* (Eoc.-Ala.), 21(1): 15, 3, 29; (Eoc.-Ga.), 21(3): 61, 10, 8; (Eoc.-Miss.), 21(4): 92, 15, 2; (Olig.-Miss.), 24(1): 8; (Eoc.-Miss.), 24(2): 32, 6, 1; *preadvenum* (Eoc.-Miss.), 24(2): 32, 6, 4; *sloani*, var. *nitida* (Rec.Atl.), 23(4): 90, 19, 9; *thalmanni* (Eoc.-Holland), 22(4): 121; *tisburyensis* (Rec.-Atl.), 24(1): 22, text figs. 1-3; *whitsettense* (Eoc.-Ga.), 21(3): 62; var. *insuetum* (Eoc.-Ga.), 21(1): 5, 1, 17-19.

Nonionella atlantica (Rec.-Atl.), 23(4): 90, 20, 4, 5; *crassipunctata* (Olig.-Miss.), 22(3): 90, 15, 16; *ecuadorana* (Mioc.-Ecuador), 24(3): 54, 9, 15; *hantkeni* (Eoc.-Ga.), 21(1): 6, 2, 1; (Eoc.-Ala.), 21(1): 15, 3, 32, 33; var. *byramensis* (Olig.-Miss.), 22(3): 89, 15, 11, 12; var. *fayettei* (Eoc.-Miss.), 21(4): 92, 15, 6; var. *spissa* (Eoc.-Ga.), 21(3): 63, 10, 12; (Eoc.-Miss.), 21(4): 92, 15, 5; (Olig.-Miss.), 22(3): 89; *jacksonensis* (Eoc.-Ala.), 21(1): 16; (Eoc.-Miss.), 21(4): 93, 15, 8; 24(1): 10; var.

compressa (Eoc.-Miss.), 21(4) : 93, 15, 7; (Olig.-Miss.), 24(1) : 8; *mauricensis* (Eoc.-Ala.), 21(1) : 16, 3, 34, 35; *oligocenica* (Olig.-Miss.), 22(3) : 89, 15, 15; **opima** (Rec.-Atl.), 23(4) : 90, 20, 1-3; *pauciloba* (Olig.-Miss.), 22(3) : 89, 15, 17; *tatumi* (Olig.-Miss.), 22(3) : 90, 15, 13, 14; 24(1) : 8.

Note on the Occurrence of *Uvigerina mexicana* Nuttall in the Ste. Croix Formation of Trinidad, B. W. I., 24(1) : 23.

Nummuloculina intermedia (Tert.-Australia), 22(1) : 15.

Nummulites antillea, 25(4) : 100; *cretacea*, 22(4) : 129; *cretaceus*, 22(4) : 129; *suprajurensis* (Jur.-Poland), 25(1) : 1.

Nummulites (*Nummulites*) *antillea*, 25(4) : 100.

[Oligocene]. A Foraminiferal Fauna from the Byram Marl at Its Type Locality, 22(3) : 76. Foraminifera from the Red Bluff-Yazoo Section at Red Bluff, Mississippi, 24(1) : 1. Note on the Occurrence of *Uvigerina mexicana* Nuttall in the Ste. Croix Formation of Trinidad, B. W. I., 24(1) : 23. *Pavoninoides*, A New Genus of the Miliolidae from Panama, 25(3) : 58.

Olympia, 22(4) : 131.

Operculinoides ellisorae (Olig.-Miss.), 22(3) : 90, 15, 20.

Ophthalmidiidae, *Polysegmentina*, A New Genus of the, 22(1) : 1.

Ophthalmidium inconstans (Rec.-Pac., Atl.), 22(1) : 15.

Orbignyna pyrei (Cret.-Venezuela), 23(1) : 8, 2, 19, 20.

[Pacific]. Three New Names for Recent Pacific Foraminifera, 24(3) : 76.

[Paleocene]. A Foraminiferal Fauna from the Paleocene of Arkansas, 22(2) : 45. *Ammobaculites paleocenicus* Cushman, A New Name, 23(4) : 77. Some New Foraminifera from the Paleocene of the Southern United States, 23(4) : 81. Additional New Foraminifera from the American Paleocene, 24(2) : 43. Some Paleocene Foraminifera from the Madruga Formation of Cuba, 24(3) : 68. Additional Species of Paleocene Foraminifera from the Madruga Formation of Cuba, 24(4) : 85.

Palmula budensis (Paleoc.-Ark.), 22(2) : 55, 9, 21, 22; *decorata*, 25(4) : 102; *delicatissima* (Paleoc.-Cuba), 24(4) : 89; *deslongchampsii* (Jur.-Poland), 25(1) : 10, 3, 1, 2; *elegantissima*, 25(4) : 102; *jarvisi* (Cret.-Trinidad), 23(2) : 43; *primitiva*, var. **paleocenica** (Paleoc.-Ala., Miss.), 24(2) : 43, 8, 7; *semi-reticulata* (Cret.-Venezuela), 23(1) : 13, 4, 5; (Cret.-Trinidad), 23(2) : 42.

- [Panama]. *Pavoninoides*, A New Genus of the Miliolidae from Panama, 25(3) : 58.
- Parrella*, and *Alabamina*, Some Notes on the Genera *Pulvinulina*, 24(1) : 13.
- Parrella*, *Pseudoparrella*, A New Generic Name, and A New Species of, 24(3) : 49.
- Parrella*, 24(1) : 13; **algeriana** (Eoc. & Olig.-Africa), 24(3) : 49, 9, 1, 2; *mexicana* (Eoc.-Peru), 25(4) : 82, 14, 18.
- Patellina advena* (Eoc.-Ga.), 21(3) : 67, 11, 1; (Olig.-Miss.), 24(1) : 9; *corrugata* (Plioc.-Alaska), 23(3) : 67, 16, 9; sp. (Eoc.-Ala.), 21(1) : 18.
- Pavonina*, 21(2) : 45; *advena*, 21(2) : 49; *agglutinans*, 21(2) : 49; *atlantica* (Rec.-Atl.), 21(2) : 48, 8, 11, 12; *flabelliformis* (Rec.-Indo-Pac.), 21(2) : 48, 8, 6-10; *italica*, 21(2) : 49; *liburnica*, 21(2) : 49; var. *trilobata* 21(2) : 49; *mexicana*, 21(2) : 49; *miocenica* (Mioc.-Fla., Jamaica), 21(2) : 46, 8, 1; *triformis* (Olig., Mioc.-Australia), 21(2) : 46, 8, 2-5.
- Pavoninoides*, A New Genus of the Miliolidae from Panama, 25(3) : 58.
- Pavoninoides**, 25(3) : 58; **panamensis** (Olig.-Panama), 25(3) : 58, text fig.
- Pelosina complanata* (Cret.-Venezuela), 23(1) : 3, 1, 2; (Cret.-Trinidad), 23(2) : 34.
- [Peru]. A New Species of *Bulimina* from the Eocene, Chira Shale, of Peru, 23(3) : 57. Foraminifera from the Eocene, Chacra Formation, of Peru, 25(3) : 49. Foraminifera from the Eocene, Verdun Formation, of Peru, 25(4) : 73.
- Planispirina contraria* (Rec.-Atl.), 22(1) : 15; *exigua* (Rec.-Pac.), 22(1) : 14.
- Planomalina almadenensis* (Cret.-Calif.), 24(4) : 98, 16, 25.
- Planorbulinella larvata* (Eoc.-Ga.), 21(3) : 73; (Olig.-Miss.), 22(3) : 102.
- Planularia arbenzi* (Mioc.-Ecuador), 24(3) : 52, 9, 6; *californica* (Plioc.-Alaska), 23(3) : 62, 15, 4-7; *charapotoa* (Mioc.-Ecuador), 24(3) : 52, 9, 9; *danvillensis* (Eoc.-Miss.), 21(4) : 85, 13, 26, 27; *deadericki* (Cret.-Ark.), 23(3) : 56, 13, 11-13; **ecuadorana** (Mioc.-Ecuador), 24(3) : 52, 9, 7, 8; **georgiana** (Eoc.-Ga.), 21(3) : 57, 9, 7; *winniana* (Eoc.-Miss.), 24(2) : 29, 5, 16, 17; sp. (Eoc.-Miss.), 21(4) : 85, 13, 28.
- Planulina alaskensis* (Plioc.-Alaska), 23(3) : 71, 16, 25, 26; **charapotoensis** (Mioc.-Ecuador), 24(3) : 67, 10, 28; *cocoaensis* (Eoc.-

- Miss.), 24(1) : 11; *ecuadorana* (Mioc.-Ecuador), 24(3) : 67, 10, 29; *limbata*, 25(4) : 101; *scanica*, 25(4) : 101.
- Plectina trinitatensis* (Eoc.-Peru), 25(4) : 77; 13, 25; *watersi* (Cret.-Venezuela), 23(1) : 8, 2, 18.
- Plectofrondicularia californica* (Mioc.-Ecuador), 24(3) : 55, 9, 16, 17; *dentifera* (Eoc.-Ecuador), 23(4) : 77, 17, 2, 3; *mexicana* (Eoc.-Miss.), 24(1) : 10; *miocenica* (Mioc.-Ecuador), 24(3) : 55, 9, 18.
- Pleurostomella clavata* (Cret.-Trinidad), 23(2) : 47; *paleocenica* (Paleoc.-Tex., Miss.), 23(4) : 86, 18, 14, 15; *subnodosa* (Cret.-Trinidad), 23(2) : 47, 12, 7; *torta* (Cret.-Trinidad), 23(2) : 47.
- [Pliocene]. Some New Species and Varieties of Foraminifera from the Pliocene of Timms Point, California, 22(2) : 65. Some New Species of Foraminifera from the Lower Pliocene of the Netherlands, 23(3) : 57. A Foraminiferal Fauna from Amchitka Island, Alaska, 23(3) : 60.
- [Poland]. Upper Jurassic Foraminifera from the Nizniow Limestone of Podole, Poland, 25(1) : 1.
- Polymorphina advena* (Eoc.-Ga.), 21(3) : 61, 10, 5; (Olig.-Miss.), 22(3) : 87, 15, 6, 7; 24(1) : 8; var. *nuda* (Eoc.-Miss.), 21(4) : 91, 14, 21; *cushmani* (Paleoc.-Ark.), 22(2) : 57, 10, 10; *fiski* (Eoc.-Miss.), 24(2) : 31, 5, 32; *frondea* (Eoc.-Ga.), 21(1) : 4, 1, 13; (Eoc.-Miss.), 21(4) : 91, 14, 20; (Olig.-Miss.), 22(3) : 88; *kincaidi* (Plioc.-Alaska), 23(3) : 64, 15, 16, 17; sp. (Eoc.-Ga.), 21(1) : 5, 1, 14; sp. A (Paleoc.-Ark.), 22(2) : 57, 10, 11.
- Polysegmentina*, A New Genus of the Ophthalmidiidae, 22(1) : 1.
- Polysegmentina** (Rec.-Pac.), 22(1) : 1; *circinata* (Rec.-Pac.), 22(1) : 1, 1, 1-4; 15.
- Proteonina testacea* from the Lower Cretaceous of the Netherlands, A New Name for, 23(3) : 72.
- Proteonina compressa* (Rec.-Pac.), 24(3) : 76; *diffflugiformis* (Eoc.-Ala.), 21(1) : 11, 3, 1; var. *calcareo* (Rec.-Atl.), 23(4) : 86, 18, 16; *eocenica* (Eoc.-Ga.), 21(1) : 1, 1, 1, 2; *hancocki* (Rec.-Pac., 24(3) : 76; *prolixa* (Eoc.-Ga.), 21(1) : 2, 1, 3; *reinholdi* (Cret.-Holland), 23(3) : 73.
- Pseudobulimina glaessneri* (Eoc.-Ala.), 21(1) : 20, 4, 19, 20.
- Pseudoclavulina* from the Upper Cretaceous of Alabama, A New, 21(3) : 74.
- Pseudoclavulina almadenensis* (Cret.-Calif.), 24(4) : 93, 16, 7; *amorpha* (Cret.-Trinidad), 23(2) : 38; *californica* (Cret.-Calif.), 24(4) : 92, 16, 6; *cocoaensis* (Olig.-Miss.), 24(1) : 5, 1, 1; *coria*

- (Cret.-Calif.), 24(4) : 93, 16, 8; **curta** (Rec.-Trinidad), 24(2) : 41, 7, 19, 20; **gracilis** (Rec.-Trinidad), 24(2) : 40, 7, 17, 18; **moorevillensis** (Cret.-Ala.), 21(3) : 74, 12, 4, 5.
- Pseudocyclammia* **bukowiensis** (Jur.-Poland), 25(1) : 5, 2, 1, 2; **rogalai** (Jur.-Poland), 25(1) : 6, 2, 3, 4.
- Pseudoglandulina* **bistegia** (Cret.-Trinidad), 23(2) : 42; *cylindracea* (Cret.-Trinidad), 23(2) : 42; **madrugaensis** (Paleoc.-Cuba), 24(3) : 70, 11, 7; *manifesta* (Paleoc.-Ark.) 22(2) : 53, 9, 6-9; (Cret.-Trinidad), 23(2) : 42; *parallela* (Cret.-Trinidad), 23(2) : 42; *pygmaea* (Paleoc.-Ark.), 22(2) : 54, 9, 10, 11.
- Pseudoparrella*, A New Generic Name, and A New Species of *Parrella*, 24(3) : 49.
- Pseudoparrella**, 24(3) : 49; **madrugaensis** (Paleoc.-Cuba), 24(3) : 73, 12, 11-13; *subperuviana* (Mioc.-Ecuador), 24(3) : 65, 10, 24, 25; *thalmanni* (Mioc.-Ecuador), 24(3) : 66, 10, 26, 27.
- Pseudophragmina* (*Proporocyclina*) *peruviana* (Eoc.-Ecuador), 22(4) : 118.
- Pseudopolymorphina* **curta** (Plioc.-Alaska), 23(3) : 63, 15, 14; *rutila* (Olig.-Miss.), 22(3) : 87, 15, 5; sp. A (Paleoc.-Ark.), 22(2) : 57, 10, 8.
- Pseudouvirgerina* **naheolensis** (Paleoc.-Ark.), 22(2) : 59, 10, 17; (Paleoc.-Cuba), 24(4) : 89.
- Pullenia* **bulloides** (Eoc.-Peru), 25(4) : 82, 14, 21; **charapotoensis** (Mioc.-Ecuador), 24(3) : 66, 10, 18; **coryelli** (Cret.-Venezuela), 23(1) : 17; (Cret.-Trinidad), 23(2) : 49; **jarvisi** (Cret.-Trinidad), 23(2) : 50; *minuta* (Cret.-Trinidad), 23(2) : 50, 12, 8; *quinqueloba*, var. *angusta* (Paleoc.-Ark.), 22(2) : 63; (Eoc. & Olig.-Miss.), 24(1) : 11, 2, 12; **salisburyi** (Plioc.-Alaska), 23(3) : 69, 16, 24.
- Pulvinulinella*, *Parrella*, and *Alabamina*, Some Notes on the Genera, 24(1) : 13.
- Pulvinulinella*, 24(1) : 13; *atlantisae* (Eoc.-Ga.), 21(3) : 70, 11, 8; *culter*, var. **midwayana** (Paleoc.-Ark.), 22(2) : 63, 11, 12; *davillensis* (Eoc.-Ga.), 21(3) : 70, 11, 9; *florealis* (Cret.-Trinidad), 23(2) : 49; *obtusa* (Eoc.-Miss.), 21(4) : 101, 16, 7, 8; (Paleoc.-Ark.), 22(2) : 63, 11, 9, 10; *subperuviana* (Mioc.-Calif.), 24(1) : 13, 2, 16.
- Pygro* **byramensis** (Olig.-Miss.), 22(3) : 83, 14, 18, 19; 24(1) : 8; *elongata* (Plioc.-Alaska), 23(3) : 62, 15, 3; **monroei** (Olig.-Miss.), 22(3) : 84, 14, 20, 21.
- Pyrulina* **cylindroides** (Paleoc.-Ark.), 22(2) : 57, 10, 6; (Cret.-

Trinidad), 23(2) : 43; *extensa* (Paleoc.-Ark.), 22(2) : 57, 10, 7; *velascoensis* (Cret.-Trinidad), 23(2) : 44.

Quadrимorphina, Species of the Genera *Allomorphina* and, 25(3) : 59.

Quadrимorphina, 25(3) : 69; *advena* (Eoc.-Calif.), 25(3) : 71, 12, 13; *allomorphinoides* (Cret., Paleoc.-Europe, Amer.), 25(3) : 69, 12, 10-12; *monterelensis* (Cret.-France), 25(3) : 70.

Quinqueloculina agglutinata (Plioc.-Alaska), 23(3) : 61, 14, 12, 13; *byramensis* (Olig.-Miss.), 22(3) : 79, 13, 18; 22(4) : 130; **compta** (Rec.-Atl.), 23(4) : 87, 19, 2; *crassa*, var. *macerata* (Olig.-Miss.), 22(3) : 79, 14, 3, 4; *glabrata* (Olig.-Miss.), 22(3) : 78, 13, 19; *hermosa* (Eoc.-Miss.), 24(2) : 28, 5, 9; *horrida* (Rec.-Atl.), 23(4) : 88, 19, 1; *laevigata* (Mioc.-Ecuador), 24(3) : 51, 9, 3; (Eoc.-Peru), 25(4) : 78, 13, 26; *longirostra* (Eoc.-Miss.), 21(4) : 81, 13, 7; *mauricensis* (Eoc.-Ala.), 21(1) : 12, 3, 5, 6; (Eoc.-Miss.), 24(2) : 28, 5, 7, 8; var. *lisbonensis* (Eoc.-Ala.), 21(1) : 12, 3, 7, 8; (Eoc.-Miss.), 21(4) : 81, 13, 9, 10; *mcglameryae* (Olig.-Miss.), 22(3) : 80, 13, 22, 23; *monroei* (Olig.-Miss.), 22(3) : 79, 14, 1, 2; *moodyensis* (Eoc.-Miss.), 21(4) : 81, 13, 8; *naheolensis* (Paleoc.-Ala.), 23(4) : 82, 17, 14; *nitida*, 22(4) : 130; *plummerae*, var. *flectata* (Paleoc.-Ala.), 23(4) : 82, 17, 15; *sabulosa* (Rec.-Atl.), 23(4) : 87, 18, 22; *transverse-striata*, 23(4) : 93; *tuberculata* (Eoc.-Miss.), 21(4) : 81, 13, 11, 12; *venusta* (Plioc.-Alaska), 23(3) : 61, 14, 16, 17; *vicksburgensis* (Olig.-Miss.), 22(3) : 78, 13, 21; *vulgaris* (Plioc.-Alaska), 23(3) : 62, 14, 14, 15; *yeguaensis* (Eoc.-Ala.), 21(1) : 12, 3, 3, 4; (Eoc.-Miss.), 24(2) : 28, 5, 5, 6; sp. (Plioc.-Alaska), 23(3) : 62, 15, 1, 2; sp. A (Jur.-Poland), 25(1) : 9, 2, 12, 13; sp. B (Jur.-Poland), 25(1) : 9, 2, 14, 15.

Ramulina aculeata (Eoc.-Ga.), 21(3) : 61, 10, 6, 7; (Paleoc.-Ark.), 22(2) : 58, 10, 12; (Cret.-Venezuela), 23(1) : 13, 4, 7, 8; (Olig.-Miss.), 24(1) : 8.

[Recent]. New Species and Varieties of Foraminifera from off the Southeastern Coast of the United States, 23(4) : 86. Some New Genera and Species of Foraminifera from Brackish Water of Trinidad, 24(1) : 15. A New Species of *Nonion* (Foraminifera) from the Woods Hole Region, 24(1) : 21. Additional New Species of Arenaceous Foraminifera from Shallow Waters of Trinidad, 24(2) : 37. New Names for Four Homonym Species

Described in "Foraminifera in the Gullmar Fjord and the Skagerrak," 24(2) : 45. Three New Names for Recent Pacific Foraminifera, 24(3) : 76.

Reophax dentalinoides (Cret.-Trinidad), 23(2) : 35, 11, 4; sp. (Cret.-Venezuela), 23(1) : 3; sp. (Cret.-Trinidad), 23(2) : 35; sp. (Eoc.-Peru), 25(3) : 50, 9, 1, 2.

Reussella, 21(2) : 23; *aculeata* (Rec.-Pac.), 21(2) : 41, 7, 10, 11; *aguayoi* (Rec.-Atl.), 21(2) : 38, 7, 6; *aperta* (Mioc.-Europe), 21(2) : 35, 6, 13; *armata* (Rec.-Australia), 21(2) : 39, 7, 7, 8; *byramensis* (Olig.-Miss.), 22(3) : 94, 16, 4, 5; 24(1) : 9; *californica* (Cret.-Calif.), 21(2) : 25, 5, 3-5; *cognata* (Olig.-Germany), 21(2) : 31, 6, 1; *cushmani* (Cret.-Sweden, France), 21(2) : 24, 5, 1; *decorata* (Mioc.-Rec.-Australia), 21(2) : 36, 6, 15-18; *elongata* (Eoc.-France, England), 21(2) : 27, 5, 10, 11; *ensiformis* (Mioc.-Rec.-Australia), 21(2) : 35, 6, 14; *eocena* (Eoc.-Fla., S. Car., Ga., Miss.), 21(2) : 30, 5, 23; (Eoc.-Miss.), 21(4) : 99, 15, 27; *glabrata* (Plioc.-Mioc., N. Car., Fla.), 21(2) : 37, 6, 21-23; *goochi* (Eoc.-La.), 21(2) : 29, 5, 17; *insueta* (Rec.-Pac.), 21(2) : 40, 7, 9; *limbata* (Eoc.-France), 21(2) : 26, 5, 8, 9; *minima* (Cret.-Sweden), 21(2) : 25, 5, 2; *miocenica* (Mioc.-Fla.), 21(2) : 36, 6, 19, 20; *oberburgensis* (Eoc.-Germany, France), 21(2) : 26, 5, 6, 7; *obtusa* (Eoc.-France), 21(2) : 27, 5, 12-14; *oligocenica* (Olig.-Miss.), 22(3) : 94, 16, 6, 7; *perforata* (Rec.-Fiji), 21(2) : 39, 7, 4; *pulchra* (Mioc.-Rec.), 21(2) : 34, 6, 11, 12; *rectimargo* (Olig.-Miss., Ala.), 21(2) : 32, 6, 2, 3; var. *hebetata* (Olig.-Miss., Ala., Fla.), 21(2) : 32, 6, 4, 5; var. *trinitatensis* (Tert.-Trinidad), 21(2) : 33, 6, 6; *recurvata* (Eoc.-France), 21(2) : 28, 5, 19; *sculptilis* (Eoc.-Ala., Miss.), 21(2) : 29, 5, 20-22; *simplex* (Rec.-Pac.), 21(2) : 40, 7, 5; *spinosissima* (Plioc.-Rec.-Medit.), 21(2) : 38, 7, 1-3; *spinulosa* (Mioc.-Austria), 21(2) : 33, 6, 8, 9; var. *atlantica* (Rec.-Atl.), 23(4) : 91, 20, 6, 7; var. *laevigata* (Mioc.-France), 21(2) : 34, 6, 10; *subrotundata* (Eoc.-Tex., La.), 21(2) : 29, 5, 18; (Eoc.-Ga.), 21(3) : 66, 10, 26; *terquemi* (Eoc.-France), 21(2) : 28, 5, 15, 16; *yumuriana* (Olig.-Cuba), 21(2) : 33, 6, 7.

Reussellinae of the Foraminiferal Family Buliminidae, The Species of the Subfamily, 21(2) : 23.

Rhabdammina discreta (Cret.-Venezuela), 23(1) : 3, 1, 1; (Cret.-Trinidad), 23(2) : 33; (Eoc.-Peru), 25(4) : 74, 13, 2; *discreta*, var. (Cret.-Trinidad), 23(2) : 33.

Robertina angusta (Eoc.-Ga.), 21(1) : 8, 2, 5; (Olig.-Miss.),

- 24(1): 9; *moodyensis* (Eoc.-Miss.), 21(4): 94, 15, 10, 11; *ovigera* (Eoc.-Ala.), 21(1): 16, 4, 4; *plummerae* (Eoc.-Miss.), 24(2): 32, 6, 6.
- Robulina inornata*, 22(4): 130; *limbata*, 25(4): 102; *umbonata*, 25(4): 102.
- Robulus alabamensis* (Paleoc.-Ark.), 22(2): 47, 7, 8; *alato-limbatus* (Eoc.-Ala.), 21(1): 13, 3, 11; (Eoc.-Miss.), 21(4): 85, 13, 25; 24(1): 10; *alto-costatus*, 22(4): 130; 23(4): 93; *americanus* (Mioc.-Ecuador), 24(3): 51, 9, 4; *arkansasana* (Paleoc.-Ark.), 22(2): 48, 7, 14; *claratus*, 25(4): 102; *degolyeri* (Paleoc.-Ark.), 22(2): 47, 7, 13; *discrepans* (Cret.-Venezuela), 23(1): 10, 3, 3; (Cret.-Trinidad), 23(2): 40; *inornatus* (Paleoc.-Ark.), 22(2): 48, 7, 9; *insuetus* (Eoc.-Ecuador), 23(4): 77, 17, 1; *insulsus* (Paleoc.-Tex.), 23(4): 83, 18, 2, 3; *limbatus*, 25(4): 102; *limbosus*, var. *hockleyensis* (Eoc.-Miss.), 24(1): 10; *macrodiscus* (Cret.-Venezuela), 23(1): 9, 3, 1, 2; (Cret.-Trinidad), 23(2): 40; (Eoc.-Peru), 25(4): 79, 14, 1; *mexicanus*, var. *alticostatus*, 22(4): 130; *midwayensis* (Paleoc.-Ark.), 22(2): 47, 7, 7; (Eoc.-Peru), 25(3): 51, 9, 7, 8; var. *virginianus*, 25(4): 102; *multiseptae* (Eoc.-Peru), 25(3): 51, 9, 6; *occidentalis*, var. *glabratus* (Eoc.-Peru), 25(4): 78, 13, 30; *oligostegius* (Cret.-Venezuela), 23(1): 10, 3, 4, 6; *pachysuturalis*, 25(4): 102; *piluliferus* (Paleoc.-Tex.), 23(4): 83, 18, 4; *pseudocassis* (Plioc.-Calif.), 22(2): 65, 12, 2; *pseudo-costatus*, var. *inornatus*, 22(4): 130; *pseudo-mamilligerus* (Paleoc.-Ark.) 22(2): 47, 7, 10; *pseudovortex* (Eoc.-Peru), 25(4): 79, 13, 31; *sternalis* (Cret.-Trinidad), 23(2): 40; *translucidus*, 25(4): 102; *turbinatus* (Paleoc.-Ark.), 22(2): 47, 7, 11; *umbonatus*, 25(4): 102; *vicksburgensis* (Olig.-Miss.), 22(3): 84, 14, 23; 24(1): 8; (Mioc.-Ecuador), 24(3): 52, 9, 5; *virginianus*, 25(4): 102; *vitrealis*, 25(4): 102; *wilcoxensis*, var. *dissentia* (Paleoc.-Ark.), 22(2): 48, 7, 12; *williamsoni* (Cret.-Trinidad), 23(2): 40; sp. (Paleoc.-Ark.), 22(2): 48, 7, 15, 16; sp. (Eoc.-Miss.), 24(2): 29, 5, 15; sp. (Jur.-Poland), 25(1): 9, 2, 16.
- Rosalina (Globotruncana) stuarti*, 22(4): 127.
- Rosalina (Rotalia) granulosa*, 22(4): 130.
- Rosalinella globigerinoides*, 22(4): 128; *lapparenti*, 22(4): 128.
- Rosalinella (Globotruncana) lapparenti*, 22(4): 130.
- Rotalia*, A New Genus, *Cribropyrgo*, and a New Species of, 22(4): 119.
- Rotalia*, Some Cuban Foraminifera of the Genus, 23(2): 23.

Rotalia byramensis (Olig.-Miss.), 22(3): 100, 16, 23; (Eoc.-Cuba), 23(2): 26, 7, 1; *capdevilensis* (Eoc.-Cuba), 23(2): 25, 6, 1-3; *ecuadorana* (Mioc.-Ecuador), 24(3): 65, 10, 17; *granulosa*, 22(4): 130; *guantanamoensis* (Olig.-Cuba), 23(2): 28, 9, 1, 2; *havanensis* (Paleoc.-Cuba), 23(2): 24, 5, 5; *jabacoensis* (Eoc.-Cuba), 23(2): 29, 10, 1-3; *madrugaensis* (Paleoc.-Cuba), 23(2): 24, 5, 4; *mexicana* (Eoc.-Cuba), 23(2): 27, 7, 3; var. *mecatepecensis* (Eoc., Olig.-Cuba), 23(2): 27, 8, 1-3; *orientalis* (Eoc.-Cuba), 23(2): 26, 7, 2; *palmerae* (Eoc.-Cuba), 23(2): 26, 6, 5-7; *parva* (Olig.-Miss.), 22(3): 100, 16, 24, 25; *peñonensis* (Eoc.-Cuba), 23(2): 25, 6, 4; *pinarensis* (Olig.-Cuba), 23(2): 28, 9, 3; *primitiva* (Cret.-Cuba), 23(2): 23, 5, 1-3; *pseudomarginata* (Eoc.-Holland), 22(4): 123; *rolshauseni* (Rec.-Gulf of Mexico), 22(4): 119, 19, 11-13; *terquemi* (Eoc.-Holland), 22(4): 123.

Rotamorphina, 25(3): 71; 25(4): 102; *cushmani* (Cret.-Trinidad), 25(3): 71, 12, 14; 25(4): 102.

Rugidia corticata, 25(1): 21.

Rzehakina epigona, var. *lata* (Cret.-Venezuela), 23(1): 9, 2, 23; (Cret.-Trinidad), 23(2): 39; var. *minima* (Cret.-Trinidad), 23(2): 39.

Saccammia rhumbleri (Cret.-Trinidad), 23(2): 34; var. *trinitatis* (Cret.-Trinidad), 23(2): 34, 11, 2; *sphaerica* (Eoc.-Peru), 25(4): 75, 13, 4, 5.

Saccorhiza ramosa (Cret.-Trinidad), 23(2): 34.

Saracenaria acutauricularis (Olig.-Miss.), 24(1): 8, 1, 6; *cushmani*, 22(4): 130; *fragilis* (Eoc.-Miss.), 24(1): 3, 1, 17; 10, 1, 17; *hantkeni* (Eoc.-Miss.), 24(1): 10, 1, 16; *triangularis* (Cret.-Venezuela), 23(1): 10, 4, 1; (Cret.-Trinidad), 23(2): 40.

Seabrookia, 25(4): 96; *cretacica* (Cret.-Cuba), 25(4): 96, 16, 18-20; *cubana* (Olig.-Cuba), 25(4): 97, 16, 22, 23; *earlandi* (Rec.-Atl., Pac.), 25(4): 98, 16, 26, 27; *lagenoides* (Eoc.-Holland), 25(4): 97, 16, 21; *pellucida* (Rec.-Pac.), 25(4): 97, 16, 24, 25.

Sigmoilina and Its Species, The Genus, 22(2): 29.

Sigmoilina victoriensis Cushman, A New Name, 22(3): 103.

Sigmoilina, 22(2): 29; *arenaria* (Rec.-Pac.), 22(2): 38, 6, 7; *arenata* (Rec.-Atl.), 22(2): 42, 6, 28; *asperula* (Mioc.-Hungary, Australia), 22(2): 33, 5, 16-18; *australis* (Rec.-Pac.), 22(2):

- 43, 6, 34; *celata* (Plioc.-Medit.), 22(2) : 36, 5, 23-29; *chapmani* (Olig.-Australia), 22(2) : 31, 5, 7-9; *costata* (Rec.-Medit.), 22(2) : 40, 6, 15-18; *edwardsi* (Rec.-Atl., Pac.), 22(2) : 39: 6, 11-13; var. *acuta* (Rec.-Antarctic), 22(2) : 40, 6, 14; *flintii* (Rec.-Atl.), 22(2) : 44, 6, 35-39; *foliacea* (Plioc.-Sicily), 22(2) : 37, 6, 5; *formosana* (Plioc.-Formosa), 22(2) : 37, 6, 6; *herzensteini* (Rec.-Arctic), 22(2) : 40, 6, 19, 20; *inconspicua* (Eoc.-La.), 22(2) : 30, 5, 4; *lapidigera* (Plioc.-Australia), 22(2) : 45; *latissima* (Rec.-Pac.), 22(2) : 44, 6, 33; *macarovi* (Rec.-Arctic), 22(2) : 40, 6, 21; *mediterranea* (Rec.-Medit.), 22(2) : 41, 6, 24-27; *miocenica* (Mioc.-Fla.), 22(2) : 33, 5, 19-22; *obesa* (Rec.-Antarctic), 22(2) : 43, 6, 29-31; *oligocenica* (Olig.-Miss., Mex., Germany), 22(2) : 31, 5, 6; *orinocoensis* (Olig.-Venezuela), 22(2) : 30, 5, 5; *ovata* (Rec.-Medit.), 22(2) : 41, 6, 22; *papyracea* (Cret.-England), 22(2) : 29, 5, 1; *schlumbergeri* (Mioc.-Rec.), 22(2) : 36, 6, 1-4; *sidebottomi* (Rec.-Medit.), 22(2) : 45; *sigmoidea* (Rec.-Atl.), 22(2) : 38, 6, 8-10; var. *compressa* (Olig.-Australia, 22(2) : 32, 5, 10-12; 22(3) : 103; *syrtica* (Rec.-Medit.), 22(2) : 41, 6, 23; *tchokrakensis* (Mioc.-Russia), 22(2) : 34; *tenuis* (Mioc.-Rec.), 22(2) : 32, 5, 13-15; *umbonata* (Rec.-Antarctic), 22(2) : 42, 6, 32; *victoriensis* (Olig.-Australia), 22(3) : 103; sp. (Eoc.-Calif.), 22(2) : 30, 5, 2; sp. (Eoc.-Calif.), 22(2) : 30, 5, 3.
- Sigmomorphina jacksonensis* (Eoc.-Ga.), 21(1) : 4, 1, 12; 21(3) : 60, 10, 3; (Eoc.-Miss.), 21(4) : 90, 14, 17, 18; 24(2) : 31, 5, 30; var. *costifera* (Eoc.-Ala.), 21(1) : 15, 3, 26; (Eoc.-Miss.), 21(4) : 90, 14, 19; *lamarcki* (Eoc.-Miss.), 21(4) : 90, 14, 15; *mindenensis* (Eoc.-Ala.), 21(1) : 14, 3, 24, 25; *semitecta*, var. *terquemiana* (Eoc.-Ala.), 21(1) : 15; (Eoc.-Ga.), 21(3) : 61, 10, 4; (Eoc.-Miss.), 21(4) : 90, 14, 11; (Paleoc.-Ark.), 22(2) : 57; (Olig.-Miss.), 22(3) : 87; (Cret.-Trinidad), 23(2) : 44; (Eoc.-Miss.), 24(2) : 31, 5, 31; *trilocularis* (Plioc.-Alaska), 23(3) 64, 15, 15; *wilcoxensis* (Paleoc.-Ark.), 22(2) : 57, 10, 9.
- Siphogenerina solitaria* (Eoc.-Peru), 25(3) : 55, 10, 7, 8.
- Siphogenerinoides eleganta* (Paleoc.-Ark.), 22(2) : 59, 10, 18; (Paleoc.-Cuba), 24(3) : 71, 11, 9; *ewaldi* (Cret.-Trinidad), 23(2) : 44, 12, 6; *parva* (Cret.-Venezuela), 23(1) : 14, 4, 14; sp. (Cret.-Venezuela), 23(1) : 14, 4, 15.
- Siphonina advena* (Olig.-Miss.), 22(3) : 100; 24(1) : 10; *clai-bornensis* (Eoc.-Ala.), 21(1) : 19; (Eoc.-Miss.), 24(2) : 35, 6, 22, 23; *danvillensis* (Eoc.-Miss.), 21(4) : 101, 16, 5, 6; 24(1) :

11; *jacksonensis* (Eoc.-Ga.), 21(3) : 69; *prima* (Paleoc.-Ark.), 22(2) : 62, 11, 7, 8.

Some Cuban Foraminifera of the Genus *Rotalia*, 23(2) : 23.

Some Cuban Species of *Globorotalia*, 25(2) : 26.

Some Foraminifera of Woodbine Age from Texas, Mississippi, Alabama, and Georgia, 22(3) : 71.

Some Homonyms in "Foraminifera from the Middle Eocene in the Southern Part of the Netherlands Province of Limburg," 22(4) : 120.

Some New Foraminifera from the American Cretaceous, 23(3) : 53.

Some New Foraminifera from the Paleocene of the Southern United States, 23(4) : 81.

Some New Genera and Species of Foraminifera from Brackish Water of Trinidad, 24(1) : 15.

Some New Species and Varieties of Foraminifera from the Pliocene of Timms Point, California, 22(2) : 65.

Some New Species of Foraminifera from the Lower Pliocene of the Netherlands, 23(3) : 57.

Some Notes on the Genera *Pulvinulinella*, *Parrella*, and *Alabamina*, 24(1) : 13.

Some Paleocene Foraminifera from the Madruga Formation of Cuba, 24(3) : 68.

Species of *Globigerina* Described between 1839 and 1850, The, 22(1) : 15.

Species of *Globigerina* Described by d'Orbigny in 1839 from Shore Sands of Cuba, The, 21(3) : 74.

Species of the Genera *Allomorphina* and *Quadriformina*, 25(3) : 59.

Species of the Genus *Chilostomella* and Related Genera, 25(4) : 84.

Species of the Subfamily Reussellinae of the Foraminiferal Family Buliminidae, The, 21(2) : 23.

Sphaeroidina and Its Species, The Genus, 25(1) : 11.

Sphaeroidina, 25(1) : 11; *austriaca* (Eoc.-Rec.), 25(1) : 15, 3, 12-17; *bulloides* (Olig.-Rec.), 25(1) : 13, 3, 8-11; *chilostomata* (Olig., Plioc.-Amer.), 25(1) : 18, 4, 10-12; *ciperana* (Olig.-Trinidad), 25(1) : 19; *compacta* (Rec.-Atl., Pac.), 25(1) : 19, 4, 14; *compressa* (Rec.-Pac.), 25(1) : 20, 4, 15, 16; *corticata*, 25(1) : 21; *cretacea*, 25(1) : 21; *dehiscens*, 25(1) : 21; var. *immatura*, 25(1) : 21; *gemmula*; 25(1) : 21; *haueri* (Mioc.-Europe), 25(1) : 16, 4, 1, 2; *murrhyna*, 25(1) : 21; *nitida* (Rec.-Phil.), 25(1) : 20,

4, 13; *ornata* (Plioc.-Fiji), 25(1) : 20, 4, 17; *parisiensis*, 25(1) : 21; *peruviana*, 25(1) : 21; *variabilis* (Olig.-Germany; Eoc.-S. Car.), 25(1) : 17, 4, 3-9; sp., 25(1) : 21.

Sphaeroidinella dehiscens, 25(1) : 21; var. *immatura*, 25(1) : 21.

Spharella, 23(4) : 93.

Spirillina limbata, var. *bipunctata* (Olig.-Miss.), 22(3) : 95; *sel-seyensis* (Eoc.-Miss.), 24(2) : 34, 6, 14, 15; *subdecorata* (Olig.-Miss.), 22(3) : 95, 16, 10; 24(1) : 9; *vicksburgensis* (Eoc.-Ga.), 21(3) : 67; (Olig.-Miss.), 22(3) : 96, 16, 11; 24(1) : 9; *vivipara* (Eoc.-Ga.), 21(3) : 67.

Spirolina arrecta (Olig.-Miss.), 22(3) : 90, 15, 18, 19.

Spiroloculina alata (Tert.-Germany), 22(1) : 14; *atlantica* (Rec.-Atl.), 23(4) : 88, 19, 3-5; *bidentata* (Eoc.-Miss.), 21(4) : 82, 13, 19; *hyalina* (Rec.-Atl.), 22(1) : 14; *lapidigera* (Plioc.-Australia, 22(2) : 45; *occlusa* (Olig.-Miss.), 22(3) : 80, 14, 7; 24(1) : 8; *spissa* (Olig.-Miss.), 22(3) : 81, 14, 6; 24(1) : 8.

Spiroplectammina dentata (Cret.-Trinidad), 23(2) : 37, 11, 3; *excolata* (Cret.-Venezuela), 23(1) : 5, 2, 2; *grzybowskii* (Cret.-Venezuela), 23(1) : 5, 4, 12, 13; (Cret.-Trinidad), 23(2) : 37; *howei* (Olig.-Miss.), 22(3) : 77, 13, 12; 24(1) : 5; *laevis*, var. *cretosa* (Paleoc.-Ark.), 22(2) : 46, 7, 1; *mississippiensis* (Eoc.-Miss.), 21(4) : 80, 13, 1; (Olig.-Miss.), 22(3) : 76, 13, 10; var. *alabamensis* (Eoc.-Ga.), 21(3) : 56, 9, 1-3; (Eoc.-Miss.), 21(4) : 80, 13, 2; (Olig.-Miss.), 22(3) : 76, 13, 11; (Eoc.-Miss.), 24(1) : 10; *paleocenica* (Paleoc.-Ala.), 23(4) : 81, 17, 17; *semicomplanata* (Cret.-Venezuela), 23(1) : 5, 2, 1; *trinitatensis* (Eoc.-Peru), 25(4) : 76, 13, 16; *wilcoxensis* (Eoc.-Miss.), 24(2) : 27, 5, 1, 2.

Taberina, 22(4) : 131.

[Taxonomic]. The Species of the Subfamily Reussellinae of the Foraminiferal Family Buliminidae, 21(2) : 23. The Species of *Globigerina* Described by d'Orbigny in 1839 from Shore Sands of Cuba, 21(3) : 74. *Polysegmentina*, A New Genus of the Ophthalmitidae, 22(1) : 1. The Genus *Hauerina* and Its Species, 22(1) : 2. The Species of *Globigerina* Described between 1839 and 1850, 22(1) : 15. The Genus *Sigmoilina* and Its Species, 22(2) : 29. *Sigmoilina victoriensis* Cushman, A New Name, 22(3) : 103. The Genus *Ceratobulimina* and Its Species, 22(4) : 107. A New Genus, *Cribropyrgo*, and a New Species of *Rotalia*, 22(4) : 119. Two New Names in the Foraminifera, 22(4) : 120. Some

Homonyms in "Foraminifera from the Middle Eocene in the Southern Part of the Netheralands Province of Limburg," 22(4): 120. Additional Homonyms in Foraminifera Erected Since 1940, 22(4): 123. *Lingulina naheolensis* Cushman, A New Name, 23(1): 18. A New Name for *Proteonina testacea* from the Lower Cretaceous of the Netherlands, 23(3): 72. *Ammobaculites paleocenicus* Cushman, A New Name, 23(4): 77. A New Genus and Some New Species of Foraminifera from the Upper Eocene of Ecuador, 23(4): 77. Homonyms in Foraminifera Erected During 1946, 23(4): 92. *Colomia*, A New Genus from the Upper Cretaceous of Cuba, 24(1): 12. Some Notes on the Genera *Pulvinulinella*, *Parrella*, and *Alabamina*, 24(1): 13. Note on the Occurrence of *Uvigerina mexicana* Nuttall in the Ste. Croix Formation of Trinidad, B. W. I., 24(1): 23. *Globigerinelloides*, A New Genus of the Globigerinidae, 24(2): 42. New Names for Four Homonym Species Described in "Foraminifera in the Gullmar Fjord and the Skagerak," 24(2): 45. *Pseudoparrella*, A New Generic Name, and A New Species of *Parrella*, 24(3): 49. Three New Names for Recent Pacific Foraminifera, 24(3): 76. *Anomalina tenuissima* (Reuss), var. *evoluta* van Bellen, A Homonym, 24(3): 76. The Genus *Coleites* and Its Species, 24(4): 81. Amendment to the Diagnosis of *Lockhartia huntii* Ovey, 24(4): 99. The Genus *Sphaeroidina* and Its Species, 25(1): 11. Some Cuban Species of *Globorotalia*, 25(2): 26. *Pavoninoides*, A New Genus of the Miliolidae from Panama, 25(3): 58. Species of the Genera *Allomorphina* and *Quadrिमorphina*, 25(3): 59. Species of the Genus *Chilostomella* and Related Genera, 25(4): 84. Homonyms in Foraminifera Erected During 1948 and 1949, 25(4): 100.

Technitella atlantica (Rec.-Atl.), 23(4): 87, 18, 17.

Terquemia, 22(4): 122.

[Texas]. Some Foraminifera of Woodbine Age from Texas, Mississippi, Alabama, and Georgia, 22(3): 71.

Textularia adalta (Eoc.-Ga.), 21(3): 57; (Eoc.-Miss.), 21(4): 80, 13, 3, 4; (Eoc.-Peru), 25(4): 76, 13, 17; *almadenensis* (Cret.-Calif.), 24(4): 91, 16, 4; *claibornensis* (Eoc.-Miss.), 24(2): 27, 5, 3, 4; *conoidea*, 22(4): 131; *conuloidea*, 22(4): 131; *cuyleri* (Eoc.-Ga.), 21(3): 57, 9, 6; *decrescens* (Plioc.-Holland), 23(3): 58, 14, 5; *dibollensis* (Eoc.-Ga.), 21(3): 56, 9, 4; (Eoc.-Miss.), 21(4): 80, 13, 5; *hannai* (Eoc.-Ga.), 21(3): 57, 9, 5; *gracillima*

- (Rec.-Atl.), 24(2): 46; *kattegatensis* (Rec.-Atl.), 24(2): 46; *plummerae*, var. (Paleoc.-Ark.), 22(2): 46, 7, 2; var. *arkansasana* (Paleoc.-Ark.), 23(4): 81, 17, 18, 19; *portenta* (Paleoc.-Ark.), 23(4): 81, 17, 16; *recta* (Olig.-Miss.), 24(1): 5; *sculpturata* (Plioc.-Holland), 23(3): 58, 14, 4; *subhauerii* (Eoc.-Ga.), 21(3): 57; (Olig.-Miss.), 22(3): 77, 13, 14; *tumidula* (Olig.-Miss.), 22(3): 77, 13, 13.
- Textulariella trinitatensis* (Cret.-Venezuela), 23(1): 8, 2, 11, 12, 17; (Cret.-Trinidad), 23(2): 39; var. *subcylindrica* (Cret.-Venezuela), 23(1): 8, 2, 10; (Cret.-Trinidad), 23(2): 39.
- Three New Names for Recent Pacific Foraminifera, 24(3): 76.
- Thurammina faerleensis* (Rec.-Atl.), 24(2): 45; *sphaerica* (Rec.-Atl.), 24(2): 45.
- Triloculina byramensis* (Olig.-Miss.), 22(3): 83, 14, 16, 17; *garretti* (Eoc.-Ala.), 21(1): 12, 3, 9; (Eoc.-Miss.), 24(2): 28, 5, 13, 14; *inuitata* (Eoc.-Miss.), 24(2): 28, 5, 11, 12; *mississippiensis* (Olig.-Miss.), 22(3): 83, 14, 15; *paulocostata* (Eoc.-Ala.), 21(1): 12, 3, 10; *pinguis*, 25(4): 102; *sculpturata* (Olig.-Miss.), 22(3): 83, 14, 14; *subrotunda*, 25(4): 102; *tricarinata* (Eoc.-Miss.), 21(4): 83, 13, 15; *trigonula* (Eoc.-Miss.), 21(4): 84, 13, 16; (Olig.-Miss.), 22(3): 83.
- Triloculina (Quinqueloculina) nitida*, 22(4): 130.
- Trimosina*, 21(2): 41; *milletti* (Rec.-Pac.), 21(2): 42, 7, 13; *orientalis* (Rec.-Pac.), 21(2): 42, 7, 14; *spinulosa* (Rec.-Pac.), 21(2): 41, 7, 12.
- [Trinidad]. Further Notes on the Cretaceous Foraminifera of Trinidad, 23(2): 31. Some New Genera and Species of Foraminifera from Brackish Water of Trinidad, 24(1): 15. Note on the Occurrence of *Uvigerina mexicana* Nuttall in the Ste. Croix Formation of Trinidad, B. W. I., 24(1): 23. Additional New Species of Arenaceous Foraminifera from Shallow Waters of Trinidad, 24(2): 37.
- Tritaxilina almadenensis* (Cret.-Calif.), 24(4): 94, 16, 9; *pleionensis* (Eoc.-Peru), 25(4): 77, 13, 22, 23.
- Trochammina comprimata* (Rec.-Trinidad), 24(2): 41, 8, 1-3; *exigua* (Cret.-Tex., Miss.), 22(3): 75, 13, 8; *globigeriniformis* (Cret.-Venezuela), 23(1): 9, 2, 21, 22; var. *altiformis* (Cret.-Trinidad), 23(2): 40; *laevigata* (Rec.-Trinidad), 24(2): 41, 7, 21, 22; *pacifica*, var. *simplex* (Rec.-Pac.), 24(3): 76, var. *simpplissima* (Rec.-Pac.), 24(3): 76; *parellela* (Cret.-Tex.), 23(3): 54, 13, 5; *pusilla* (Rec.-Atl.), 24(2): 46; *quadriloba*

(Rec.-Atl.), 24(2): 46; *rainwateri* (Cret.-Tex., Miss., Ala.), 22(3): 75, 13, 9; *teasi* (Eoc.-Peru), 25(3): 51, 9, 5; 25(4): 78, 13, 27, 28; *texana* (Cret.-Trinidad), 23(2): 40, 11, 5.
Trochamminita, 24(1): 17; *irregularis* (Rec.-Trinidad), 24(1): 17, 4, 1-3.

Trochamminoides velascoensis (Cret.-Trinidad), 23(2): 36.

Trocholina floridana (Cret.-Fla.), 23(2): 29, 10, 6-10.

Truncorotalia (subgenus of *Globorotalia*), 25(2): 35.

Turborotalia (subgenus of *Globorotalia*), 25(2): 42.

Two New Names in the Foraminifera, 22(4): 120.

Two New Species of Foraminifera from the Arkadelphia Marl of Arkansas, 23(3): 56.

Two New Species of Lower Cretaceous Foraminifera from Florida, 23(2): 29.

Upper Jurassic Foraminifera from the Nizniow Limestone of Podole, Poland, 25(1): 1.

Uvigerina mexicana Nuttall in the Ste. Croix Formation of Trinidad, B. W. I., Note on the Occurrence of, 24(1): 23.

Uvigerina beccarii (Olig.-Trinidad), 24(1): 23; *charapotoensis* (Mioc.-Ecuador), 24(3): 62, 10, 4; *cushmani* (Plioc.-Alaska), 23(3): 66, 16, 4, 5; *farinosa* (Eoc.-Peru), 25(4): 80, 14, 9; *gardnerae*, var. *texana* (Eoc.-Miss.), 24(1): 10; *garzaensis* (Eoc.-Peru), 25(4): 80, 14, 10; *jacksonensis* (Eoc.-Peru), 25(4): 80, 14, 11; *marksii* (Mioc.-Ecuador), 24(3): 62, 10, 6; *mexicana* (Olig.-Trinidad), 24(1): 23; *minuta* (Eoc.-Peru), 25(3): 54, 10, 5, 6; *vicksburgensis* (Olig.-Miss.), 24(1): 9, 1, 9; *yazooensis* (Eoc.-Miss.), 24(1): 10, 2, 7.

Uvigerinella californica (Mioc.-Ecuador), 24(3): 61, 10, 7; *obesa* (Mioc.-Ecuador), 24(3): 61, 10, 5.

Vaginulina cretacea (Cret.-Venezuela), 23(1): 12, 4, 2; *longiforma* (Paleoc.-Ark.), 22(2): 54, 9, 13, 14; (Paleoc.-Cuba), 24(3): 70, 11, 4; *plumoides* (Paleoc.-Ark.), 22(2): 55, 9, 20; *robusta* (Paleoc.-Ark.), 22(2): 54, 9, 15-19; *semilaevis* (Paleoc.-Cuba), 24(3): 70, 11, 6; *striatula*, 22(4): 129; *subbadensis* (Plioc.-Calif.), 22(2): 66, 12, 6, 7; var. *aculeata* (Plioc.-Calif.), 22(2): 67, 12, 8-10; var. *serrulata* (Plioc.-Calif.), 22(2): 67, 12, 11; sp. A (Paleoc.-Ark.), 22(2): 55, 8, 11.

Valvulamina nassauensis, var. *cubana* (Paleoc.-Cuba), 24(3): 68, 11, 1.

Valvulineria, 25(3) : 71; 25(4) : 102; *allomorphinoides* (Paleoc.-Ark.), 22(2) : 62, 11, 1; (Cret.-Trinidad), 23(2) : 48; 25(3) : 71, 12, 14; *californica* (Mioc.-Calif.), 25(3) : 72, 12, 15; *casitasensis* (Mioc.-Ecuador), 24(3) : 63, 10, 9; *chirana* (Mioc.-Ecuador), 24(3) : 64, 10, 11; *cushmani*, 25(4) : 102; *ecuadorana* (Mioc.-Ecuador), 24(3) : 64, 10, 12; *extensa* (Paleoc.-Cuba), 24(3) : 72, 12, 7-9; *georgiana*, var. *compacta* (Eoc.-Ga.), 21(1) : 9, 2, 10; *insueta* (Paleoc.-Cuba), 24(4) : 85, 14, 7-9; *jacksonensis*, var. *dentata* (Eoc.-Ga.), 21(1) : 9, 2, 11; *lobata* (Cret.-Miss.), 23(3) : 55, 13, 6; *madrugaensis* (Paleoc.-Cuba), 24(3) : 73, 12, 10; *malagaensis* (Mioc.-Ecuador), 24(3) : 63, 10, 10; *paucilocula* (Olig.-Miss.), 22(3) : 98, 16, 16; *suturalis* (Eoc.-Peru), 25(3) : 55, 10, 10, 11; *texana* (Eoc.-Ga.), 21(3) : 68, 11, 6.

[Venezuela]. A Foraminiferal Fauna from the Santa Anita Formation of Venezuela, 23(1) : 1.

Ventilabrella carseyae (Cret.-Venezuela), 23(1) : 14, 4, 9, 10.

Vermiculum (*Triloculina*) *subrotundum*, 25(4) : 102.

Verneuilina anglica (Jur.-Poland), 25(1) : 6, 1, 13; *liasina* (Jur.-Poland), 25(1) : 6, 1, 11, 12; *polonica* (Jur.-Poland), 25(1) : 7, 1, 14, 15.

Virgulina californiensis (Mioc.-Ecuador), 24(3) : 58, 9, 27, 28; *dibollensis* (Eoc.-Ala.), 21(1) : 17, 4, 7; (Eoc.-Ga.), 21(3) : 65, 10, 19; (Eoc.-Miss.), 21(4) : 95, 15, 13; *dibollensis*, var. (Eoc.-Miss., 24(2) : 33, 6, 8; *diversa* (Eoc.-Peru), 25(3) : 53, 9, 17; *jurassica* (Jur.-Poland), 25(1) : 10, 3, 4, 5; *mcguirti* (Eoc.-Ga.), 21(3) : 65, 10, 22; *minutissima* (Eoc.-Ga.), 21(1) : 8, 2, 7; *navarroana* (Cret.-Trinidad), 23(2) : 46; *nodosa* (Plioc.-Alaska), 23(3) : 66, 16, 1; *recta* (Eoc.-Ga.), 21(3) : 65, 10, 18; *restinensis* (Eoc.-Peru), 25(3) : 53, 9, 15, 16; *unca* (Eoc.-Miss.), 24(1) : 4, 2, 3; 10, 2, 3; *vicksburgensis* (Olig.-Miss.), 22(3) : 92, 15, 28; 24(1) : 9; *yazooensis* (Eoc.-Miss.), 24(1) : 3, 2, 4; 10, 2, 4; *zetina* (Eoc.-Ala.), 21(1) : 17, 4, 8; (Eoc.-Ga.), 21(3) : 65, 10, 20, 21.

Vulvulina jarvisi (Eoc.-Peru), 25(4) : 77, 13, 18, 19.